



# SPIRIT 2025 statement: updated guideline for protocols of randomised trials

An-Wen Chan, <sup>1</sup> Isabelle Boutron, <sup>2,3</sup> Sally Hopewell, <sup>4</sup> David Moher, <sup>5</sup> Kenneth F Schulz, <sup>6</sup> Gary S Collins, <sup>7</sup> Ruth Tunn, <sup>4</sup> Rakesh Aggarwal, <sup>8</sup> Michael Berkwits, <sup>9</sup> Jesse A Berlin, <sup>10,11</sup> Nita Bhandari, <sup>12</sup> Nancy J Butcher, <sup>13,14</sup> Marion K Campbell, <sup>15</sup> Runcie C W Chidebe, <sup>16,17</sup> Diana R Elbourne, <sup>18</sup> Andrew J Farmer, <sup>19</sup> Dean A Fergusson, <sup>20</sup> Robert M Golub, <sup>21</sup> Steven N Goodman, <sup>22</sup> Tammy C Hoffmann, <sup>23</sup> John P A Ioannidis, <sup>24</sup> Brennan C Kahan, <sup>25</sup> Rachel L Knowles, <sup>26</sup> Sarah E Lamb, <sup>27</sup> Steff Lewis, <sup>28</sup> Elizabeth Loder, <sup>29,30</sup> Martin Offringa, <sup>13</sup> Philippe Ravaud, <sup>31</sup> Dawn P Richards, <sup>32</sup> Frank W Rockhold, <sup>33</sup> David L Schriger, <sup>34</sup> Nandi L Siegfried, <sup>35</sup> Sophie Staniszewska, <sup>36</sup> Rod S Taylor, <sup>37</sup> Lehana Thabane, <sup>38,39</sup> David J Torgerson, <sup>40</sup> Sunita Vohra, <sup>41</sup> Ian R White, <sup>25</sup> Asbjørn Hróbjartsson <sup>42,43</sup>

For numbered affiliations see end of the article

Correspondence to: A-W Chan anwen.chan@utoronto.ca; (ORCID 0000-0002-4498-3382)

Additional material is published online only. To view please visit the journal online.

Cite this as: *BMJ* 2025;389:e081477 http://dx.doi.org/10.1136/ bmj-2024-081477

Accepted: 31 January 2025

## **ABSTRACT**

#### **IMPORTANCE**

The protocol of a randomised trial is the foundation for study planning, conduct, reporting, and external review. However, trial protocols vary in their completeness and often do not address key elements of design and conduct. The SPIRIT (Standard Protocol Items: Recommendations for Interventional Trials) statement was first published in 2013 as guidance to improve the completeness of trial protocols. Periodic updates incorporating the latest evidence and best practices are needed to ensure that the guidance remains relevant to users.

## OBJECTIVE

To systematically update the SPIRIT recommendations for minimum items to address in the protocol of a randomised trial.

# DESIGN

We completed a scoping review and developed a project specific database of empirical and theoretical evidence to generate a list of potential changes to the SPIRIT 2013 checklist. The list was enriched with recommendations provided by lead authors of existing SPIRIT/CONSORT (Consolidated Standards of Reporting Trials) extensions (Harms, Outcomes, Non-pharmacological Treatment) and other reporting guidelines (TiDieR). The potential modifications were rated in a three-round Delphi survey followed by a consensus meeting.

# **SUMMARY POINTS**

A complete, transparent, and accessible protocol is critical for the planning, conduct, reporting, and external review of randomised trials

The SPIRIT (Standard Protocol Items: Recommendations for Interventional Trials) 2025 statement provides guidance on items to address in trial protocols, reflecting methodological advances and feedback from users

The SPIRIT 2025 statement consists of a checklist of 34 minimum items; a diagram illustrating the schedule of enrolment, interventions, and assessments; and an expanded checklist that details the critical elements of each checklist item

Research teams, sponsors, editors, funders, and research ethics committees/institutional review boards should use and endorse SPIRIT 2025 to promote transparent protocol content

## **FINDINGS**

Overall, 317 individuals participated in the Delphi consensus process and 30 experts attended the consensus meeting. The process led to the addition of two new protocol items, revision to five items, deletion/merger of five items, and integration of key items from other relevant reporting guidelines. Notable changes include a new open science section, additional emphasis on the assessment of harms and description of interventions and comparators, and a new item on how patients and the public will be involved in trial design, conduct, and reporting. The updated SPIRIT 2025 statement consists of an evidence based checklist of 34 minimum items to address in a trial protocol, along with a diagram illustrating the schedule of enrolment, interventions, and assessments for trial participants. To facilitate implementation, we also developed an expanded version of the SPIRIT 2025 checklist and an accompanying explanation and elaboration document.

## **CONCLUSIONS AND RELEVANCE**

Widespread endorsement and adherence to the updated SPIRIT 2025 statement have the potential to enhance the transparency and completeness of trial protocols for the benefit of investigators, trial participants, patients, funders, research ethics committees, journals, trial registries, policymakers, regulators, and other reviewers.

## Introduction

"Readers should not have to infer what was probably done; they should be told explicitly." Douglas G Altman<sup>1</sup>

Robustly designed, properly conducted, and fully reported randomised trials underpin evidence based practice and policy. As the most important record of planned methods and conduct, a well written protocol has a key role in promoting consistent and rigorous execution by the trial team. The protocol also serves as the basis for oversight and review of scientific, ethical, safety, and operational issues by funders, regulators, research ethics committees/institutional review boards (REC/IRB), journal editors, researchers, patients, and the public.<sup>2-9</sup> After trial completion, the protocol is essential for understanding and interpreting the results.

Despite the central role of protocols, there is substantial variation in the completeness of protocol content. <sup>10</sup> <sup>11</sup> Many trial protocols do not adequately describe important elements including the primary outcomes, treatment allocation methods, use of blinding, measurement of adverse events, sample size calculations, data analysis methods, dissemination policies, and roles of sponsors and investigators in trial design. <sup>10-12</sup> Gaps in protocol content can lead to avoidable protocol amendments, <sup>13</sup> inconsistent or poor trial conduct, and lack of transparency in terms of what was planned and implemented.

In response to these protocol deficiencies, the SPIRIT (Standard Protocol Items: Recommendations for Interventional Trials) guidance was first published in 2013. 14 15 Aligned with the CONSORT (Consolidated Standards of Reporting Trials) guidance for reporting completed trials, 16 the international SPIRIT initiative aims to improve the completeness of trial protocols by producing evidence based recommendations for a minimum set of items to be addressed in protocols. The SPIRIT 2013 guidance has been translated into seven languages and is widely endorsed by national funders, research organisations, over 150 medical journals, and the World Association of Medical Editors.

In January 2020, the SPIRIT and CONSORT executive groups held a joint meeting in Oxford, UK, to discuss strategic planning. There was broad recognition of the need to update both checklists to reflect the evolving trials environment and methodological advancements, including the growing international support for improved research transparency, accessibility, and reproducibility (collectively referred to as open science)<sup>17</sup> as well as greater patient and public involvement in research.

As the SPIRIT 2013 and CONSORT 2010 statements were conceptually linked with overlapping content and implementation strategies, the two groups decided to merge into the joint SPIRIT-CONSORT executive group and to update both checklists simultaneously. The joint update was an opportunity to further align the checklists and provide consistent guidance in the reporting of trial design, conduct, and analysis—from study conception to the publication of results. Harmonising the reporting recommendations could help improve usability and adherence. Here, we introduce the updated SPIRIT 2025 statement; the CONSORT 2025 statement is published separately.

# Methods

The methods have been detailed elsewhere. <sup>19</sup> <sup>20</sup> In brief, we followed the EQUATOR Network guidance for developers of health research guidelines. <sup>21</sup> We first conducted a scoping review of the literature from 2013 to 2022 to identify published comments suggesting modifications or reflecting on the strengths and challenges of SPIRIT 2013; these findings have been published separately. <sup>22</sup> We also conducted a broader search for empirical and theoretical evidence published from 2013 to 2024 that was relevant to SPIRIT and risk of bias in randomised

trials, producing the SPIRIT-CONSORT Evidence Bibliographic database.<sup>23</sup> The evidence identified in the literature was combined with recommendations provided by the lead authors of key SPIRIT and CONSORT extensions (Harms,<sup>24</sup> Outcomes,<sup>25</sup> Non-pharmacological Treatment<sup>26</sup>), and the Template for Intervention Description and Replication (TIDieR),<sup>27</sup> along with user feedback.

Based on the gathered evidence, a preliminary list of five potential additions to the SPIRIT 2013 checklist was created for review in an international, three-round online Delphi survey. A total of 317 participants were recruited through professional research networks, societies, and the project website. Participants represented a broad range of roles in clinical trials, including statisticians/methodologists/ epidemiologists (n=198), trial investigators (n=73), systematic reviewers/guideline developers (n=73), clinicians (n=58), journal editors (n=47), and patients and members of the public (n=17) (numbers are not mutually exclusive). During each survey round. participants rated the importance of modifications on a five-point Likert scale and provided comments or suggestions for additional items. A high level of agreement was defined by at least 80% of respondents rating the importance of a proposed modification as high (score of 4 or 5) or low (score of 1 or 2).

The Delphi survey results were then discussed at a two-day online consensus meeting in March 2023, attended by 30 invited international experts representing a range of relevant groups. Meeting participants discussed potential new and modified SPIRIT checklist items, with anonymous polling of participants in cases of ongoing disagreement.

The executive group met in person in April 2023 to develop a draft checklist based on the consensus meeting discussion. After a further round of review by consensus meeting participants, the executive group finalised the SPIRIT 2025 statement.

## **Updated SPIRIT 2025 statement**

The SPIRIT 2025 statement comprises a checklist of 34 minimum protocol items (table 1) and a diagram illustrating the schedule of enrolment, interventions, and assessments (fig 1). An accompanying SPIRIT 2025 explanation and elaboration document provides background and context for each checklist item along with examples of good reporting. We strongly recommend that the SPIRIT 2025 explanation and elaboration document be used routinely alongside the SPIRIT 2025 statement to facilitate better understanding of and adherence to the checklist items.

To present the recommendations in diverse formats, we also developed an expanded version of the SPIRIT 2025 checklist with bullet points of key issues to consider for each item (appendix 1), as done with other initiatives.<sup>29-31</sup> The expanded checklist comprises an abridged version of elements presented in the SPIRIT 2025 explanation and elaboration document,<sup>28</sup> with examples and references removed.

Section/topic	No	SPIRIT 2025 checklist item description
Administrative informa	ation	
Fitle and structured	1a	Title stating the trial design, population, and interventions, with identification as a protocol
summary	1b	Structured summary of trial design and methods, including items from the World Health Organization Trial Registration Data Set
Protocol version	2	Version date and identifier
Roles and	3a	Names, affiliations, and roles of protocol contributors
esponsibilities .	3b	Name and contact information for the trial sponsor
•	3c	Role of trial sponsor and funders in design, conduct, analysis, and reporting of trial; including any authority over these activities
	3d	Composition, roles, and responsibilities of the coordinating site, steering committee, endpoint adjudication committee, data management team and other individuals or groups overseeing the trial, if applicable
Open science		
rial registration	4	Name of trial registry, identifying number (with URL), and date of registration. If not yet registered, name of intended registry
Protocol and statistical analysis plan		Where the trial protocol and statistical analysis plan can be accessed
Data sharing	6	Where and how the individual de-identified participant data (including data dictionary), statistical code, and any other materials will be accessible
unding and conflicts	7a	Sources of funding and other support (eg, supply of drugs)
of interest	7b	Financial and other conflicts of interest for principal investigators and steering committee members
Dissemination policy	8	Plans to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, reporting in trial registry, plain language summary, publication)
ntroduction		
Background and rationale	9a	Scientific background and rationale, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention
	9b	Explanation for choice of comparator
Objectives	10	Specific objectives related to benefits and harms
Methods: Patient and	public	involvement, trial design
Patient and public nvolvement	11	Details of, or plans for, patient or public involvement in the design, conduct, and reporting of the trial
Trial design	12	Description of trial design including type of trial (eg, parallel group, crossover), allocation ratio, and framework (eg, superiority, equivalence, non-inferiority, exploratory)
Methods: Participants	, inter	ventions, and outcomes
rial setting	13	Settings (eg, community, hospital) and locations (eg, countries, sites) where the trial will be conducted
Eligibility criteria	14a	Eligibility criteria for participants
	14b	If applicable, eligibility criteria for sites and for individuals who will deliver the interventions (eg, surgeons, physiotherapists)
Intervention and comparator	15a	Intervention and comparator with sufficient details to allow replication including how, when, and by whom they will be administered. If relevant where additional materials describing the intervention and comparator (eg, intervention manual) can be accessed
·	15b	Criteria for discontinuing or modifying allocated intervention/comparator for a trial participant (eg, drug dose change in response to harms, participant request, or improving/worsening disease)
	15c	Strategies to improve adherence to intervention/comparator protocols, if applicable, and any procedures for monitoring adherence (eg, drug tablet return, sessions attended)
•	15d	Concomitant care that is permitted or prohibited during the trial
Outcomes	16	Primary and secondary outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome
Harms	17	How harms are defined and will be assessed (eg, systematically, non-systematically)
Participant timeline	18	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see fig 1)
Sample size	19	How sample size was determined, including all assumptions supporting the sample size calculation
Recruitment	20	Strategies for achieving adequate participant enrolment to reach target sample size
Methods: Assignment	of inte	erventions
Randomisation:		
Sequence	21a	Who will generate the random allocation sequence and the method used
generation	21b	Type of randomisation (simple or restricted) and details of any factors for stratification. To reduce predictability of a random sequence, other details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions
Allocation concealment	22	Mechanism used to implement the random allocation sequence (eg, central computer/telephone; sequentially numbered, opaque, sealed containers), describing any steps to conceal the sequence until interventions are assigned
mechanism Implementation	23	Whether the personnel who will enrol and those who will assign participants to the interventions will have access to the random allocation
Blinding	24a	sequence Who will be blinded after assignment to interventions (eg, participants, care providers, outcome assessors, data analysts)
	24b	If blinded, how blinding will be achieved and description of the similarity of interventions
	24c	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial
		anagement, and analysis
Data collection methods	25a	Plans for assessment and collection of trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of trial instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known.
	25b	Reference to where data collection forms can be accessed, if not in the protocol Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who
		discontinue or deviate from intervention protocols

(Continued)

# RESEARCH METHODS AND REPORTING

Table 1   Continued			
Section/topic	No	SPIRIT 2025 checklist item description	
Statistical methods	27a	Statistical methods used to compare groups for primary and secondary outcomes, including harms	
	27b	Definition of who will be included in each analysis (eg, all randomised participants), and in which group	
	27c	How missing data will be handled in the analysis	
	27d	Methods for any additional analyses (eg, subgroup and sensitivity analyses)	
<b>Methods: Monitoring</b>			
Data monitoring committee	28a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent fro sponsor and funder; conflicts of interest and reference to where further details about its charter can be found, if not in the protocol. Alter an explanation of why a DMC is not needed	
	28b	Explanation of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	
Trial monitoring	29	Frequency and procedures for monitoring trial conduct. If there is no monitoring, give explanation	
Ethics			
Research ethics approval	30	Plans for seeking research ethics committee/institutional review board approval	
Protocol amendments	31	Plans for communicating important protocol modifications to relevant parties	
Consent or assent	32a	Who will obtain informed consent or assent from potential trial participants or authorised proxies, and how	
	32b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	
Confidentiality	33	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	
Ancillary and post-trial care	34	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	
SPIRIT=Standard Protocol	Items: R	ecommendations for Interventional Trials.	

## **Main changes**

Substantive changes made in this update are detailed in box 1. We added two new checklist items, revised the content of five items, deleted three items, merged two items, and integrated key items from CONSORT Harms 2022, <sup>24</sup> SPIRIT-Outcomes 2022, <sup>25</sup> and TIDieR<sup>27</sup> into the main checklist and explanatory document. We also restructured the SPIRIT checklist and created a new open science section consolidating items critical to promoting access to information about trial methods and results, including trial registration; sharing of the full protocol, statistical analysis plan, and de-

identified participant level data; and disclosure of funding sources and conflicts of interest. We have also harmonised the wording between SPIRIT and CONSORT checklist items and clarified the wording of some items. A comparison of the SPIRIT 2025 and 2013 checklists is available in appendix 2.

# Definition of a randomised trial protocol

The protocol is a central document that provides sufficient detail to enable (a) understanding of the rationale, objectives, population, interventions, methods, statistical analyses, ethical considerations,

	Trial period											
	Enrolment		Post-randomisation					Close -out				
Timepoint*	<i>-t<sub>i</sub></i> to 0	0	t,	<b>t</b> <sub>2</sub>	<b>t</b> <sub>3</sub>	t <sub>4</sub>	etc	t <sub>x</sub>				
Enrolment		•	•									
Eligibility screen	Х											
Informed consent	Х											
(List other procedures)	Х											
Randomisation		Χ										
Intervention or comparator												
(Intervention)†		Х —						→ X				
(Comparator)‡		Χ		Х		Х						
Assessments				•		•	•	•				
(List baseline variables and tests)	Х	Χ										
(List outcome variables and tests)			Х	Х	Х	Х	etc	Х				
(List other data variables and tests)				Х		Х	etc	Х				

Fig 1 | SPIRIT 2025 diagram of the schedule of enrolment, interventions, and assessments. Recommended content can be displayed using various schematic formats. \*List target time points and acceptable time windows in this row (eg, 30±3 days). †Arrow indicates continuous delivery of intervention (eg, drug). ‡Example illustrates delivery of comparator at discrete time points (eg, psychotherapy)

## Box 1: Summary of main changes in SPIRIT 2025

#### Addition of new checklist items:

- Item 11: added item on how patients and the public are involved in the design, conduct, and reporting of the trial
- Item 29: added item on trial monitoring (replaces prior item on auditing)

## Revised content of checklist items

- Item 4: revised item to include date of trial registration
- Item 5: revised item to include where the statistical analysis plan can be accessed in addition to the trial protocol (previously covered under item on statistical methods)
- Item 7b: revised item to include financial and other conflicts of interest of steering committee members
- Item 24a/24b: split item into separate sub-items covering (a) who will be blinded and (b) how, and revised to include description of the similarity of interventions
- Item 27d: revised to refer to sensitivity analyses

# Deletion/merger of checklist items

- Deleted item on auditing (replaced with trial monitoring)
- Deleted appendix items:
- Informed consent materials
- Biological specimens
- Merged item on access to data for trial investigators with item on data management (item 26)
- Merged item on authorship eligibility guidelines and use of professional writers with item on dissemination policy (item 8)

# Integration of checklist items from CONSORT Harms 2022 and TIDieR

• Revised items to emphasise reporting of harms (items 10, 17, 27a)<sup>24</sup> and to call for additional details relating to interventions and comparators (item 15a)<sup>27</sup>

## Structure and organisation of checklist items

- Created a new section on open science that includes trial registration (item 4), access to the trial protocol and statistical analysis plan (item 5), plans for sharing de-identified participant level data (item 6), funding and conflicts of interest (item 7), and plans for dissemination of trial results (item 8)
- Item 14a/b: split item into separate sub-items covering eligibility criteria for (a) participants and (b) sites and personnel
- Item 27b/27c: split item into separate sub-items covering the analysis population and methods for handling missing data
- Aligned wording of SPIRIT checklist items with that of CONSORT checklist items, and vice versa
- Clarified and simplified wording of some items.

CONSORT=Consolidated Standards of Reporting Trials; SPIRIT=Standard Protocol Items: Recommendations for Interventional Trials; TIDieR=Template for Intervention Description and Replication.

dissemination plans, and administration of the trial; (b) replication of trial methods and conduct; and (c) appraisal of trial validity, feasibility, and ethical rigour.<sup>14</sup>

The full protocol must be submitted for approval by an REC/IRB before enrolling participants.  $^{32}$  As a living document that is often formally amended during the trial,  $^{13}$   $^{33}$  every protocol version should contain a transparent audit trail documenting the dates and descriptions of changes. Important protocol amendments should be reported to REC/IRBs and trial registries as they occur, and subsequently described in reports of completed trials.  $^{34}$ 

# Scope of SPIRIT 2025

SPIRIT 2025 addresses the minimum content of a protocol, focusing on the most common type of randomised trial—the two-group parallel design. However, most of the SPIRIT items are relevant to any type of trial. SPIRIT 2025 has been designed to complement and enhance the expanding trial registration requirements mandated by legislation, journals, and funding policies.<sup>35</sup> SPIRIT 2025 encompasses and builds upon recommendations from the International Council for Harmonization Good Clinical Practice E6(R3) guidance<sup>36</sup> and 2024

Declaration of Helsinki,<sup>32</sup> including the Declaration of Helsinki's requirement that the protocol address potential conflicts of interest and provision of post-trial

It is feasible to address all SPIRIT 2025 checklist items in a single protocol document, as illustrated by the examples we identified from existing protocols for every item. There are often related documents (eg, full statistical analysis plan, at management plan) that provide further details on specific items. Any such documents should be referenced in the protocol and made available for review.

The main purpose of SPIRIT 2025 is to promote transparency and an adequate description of what is planned—not to prescribe how a trial should be designed or conducted. The checklist also does not focus on the protocol format, which is often subject to local regulations or practice. The checklist should not be used to appraise the quality of trial design or conduct, as it is possible for the protocol of a poorly designed trial to address all checklist items by fully describing its inadequate design and conduct features. Recent guidance from the World Health Organization (WHO) outlines best practices for designing and conducting trials.<sup>38</sup>

## **Implementation**

The SPIRIT 2025 statement supersedes the SPIRIT 2013 statement, which should no longer be used or cited. We encourage research organisations, sponsors, funders, REC/IRBs, journal editors and publishers to endorse SPIRIT 2025 and request that they update their resources and instructions to research teams and reviewers with reference to the updated guidance.

When protocols are submitted for review or publication, we recommend the submission of a completed SPIRIT 2025 checklist that indicates where (eg, page number) checklist items are reported in the protocol. Trial investigators and sponsors should address all SPIRIT 2025 checklist items in the protocol before REC/IRB submission. If an item is not relevant for a particular trial (eg, no interim analysis planned), then this should be explicitly stated along with an explanation. We encourage investigators to ensure consistency of information in the protocol, related documents (eg, full statistical analysis plan), <sup>37</sup> and trial registry record. <sup>39</sup>

To facilitate implementation, a new SPIRIT-CONSORT website (consort-spirit.org) provides resources based on the SPIRIT and CONSORT 2025 statements, including a fillable checklist, protocol writing tools, and training materials for researchers, trainees, journal editors, peer reviewers, patients, and the public.

## Limitations

As a minimum standard focused on parallel group randomised trials, SPIRIT 2025 may not encompass every protocol item relevant for a particular trial. For example, a factorial trial design has additional analytical considerations related to potential statistical interactions, 40 and trials evaluating patient reported outcomes have specific considerations regarding data collection methods. 41 Extensions to SPIRIT 2013 were developed to provide additional guidance on reporting different types of trial designs, data, and interventions. 25 34 40-47 We will engage with the leaders of these extensions to implement a process for aligning them with the updated SPIRIT 2025 statement. In the meantime, we recommend that the existing version of the relevant SPIRIT extensions be used.

## **Potential impact**

The updated SPIRIT 2025 statement and its accompanying explanation and elaboration document can be helpful in several ways. SPIRIT 2025 will continue to serve as an educational resource for new investigators, trainees, peer reviewers, and REC/IRB members. The explicit incorporation of an open science section in the SPIRIT checklist will support the growing global push for greater transparency and sharing of trial materials and outputs to facilitate evidence synthesis and reproducibility of research.

Trial investigators can consult the guidance when drafting their protocols to ensure that all elements are addressed. Meta-research reviews of protocols have found improved completeness of protocol content after

SPIRIT 2013 was introduced. <sup>10</sup> <sup>11</sup> <sup>48</sup> <sup>49</sup> In addition to improved reporting, adherence to SPIRIT 2025 may promote high quality trial design and implementation because SPIRIT is used during the planning stage of a trial. This provides an opportunity to improve the validity and successful completion of trials by reminding investigators about important issues to consider before the study begins. Better protocols can also help study personnel to implement the trial consistently across sites.

Another potential benefit of SPIRIT 2025 is its impact on administrative burden. Improved completeness of protocols may improve the efficiency of external review by reducing avoidable queries to investigators about incomplete or unclear protocol related information. 50 51 High quality protocols addressing all SPIRIT items may also help to reduce the number and burden of protocol amendments during the trial-many of which can be avoided with careful consideration of key issues when developing the protocol.13 33 Widespread adoption of SPIRIT 2025 as a common standard across REC/ IRBs, funding agencies, regulatory agencies, and journals could simplify the work of trial investigators and sponsors because a SPIRIT based protocol would then fulfil the harmonised application requirements of multiple groups.

Further, adherence to SPIRIT 2025 may help ensure that protocols contain the requisite information for critical appraisal and trial interpretation by peer reviewers, funders, REC/IRBs, and journals.7 High quality protocols provide important information about trial methods and conduct that is usually not available in trial registries or publications reporting completed trials. As a transparent record of the investigators' original intent, comparison of protocols with reports of completed trials helps to identify selective reporting of results and undisclosed amendments, such as changes to primary outcomes or analyses. 52 53 These benefits of SPIRIT based protocols can only be fully realised when trial protocols are routinely made publicly available through trial registries (eg, PDF upload), journals, and online repositories. 7 54 55

The SPIRIT 2025 statement incorporates new evidence and emerging perspectives to ensure that the guidance remains relevant to users. Widespread endorsement and adoption of the updated recommendations have the potential to improve protocol content and implementation; facilitate registration, oversight, and appraisal of trials; and ultimately enhance transparency and translation to better healthcare.

## **AUTHOR AFFILIATIONS**

<sup>1</sup>Department of Medicine, Women's College Research Institute, University of Toronto, Toronto, ON, M5S 1B2, Canada

<sup>2</sup>Université Paris Cité and Université Sorbonne Paris Nord, Inserm, INRAE, Centre for Research in Epidemiology and Statistics (CRESS), Paris, France

<sup>3</sup>Centre d'Epidémiologie Clinique, Hôpital Hôtel Dieu, AP-HP, Paris, France

<sup>4</sup>Oxford Clinical Trials Research Unit, Centre for Statistics in Medicine, University of Oxford, Oxford, UK

<sup>5</sup>Centre for Journalology, Clinical Epidemiology Programme, Ottawa Hospital Research Institute, Ottawa, ON, Canada

<sup>6</sup>Department of Obstetrics and Gynecology, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA <sup>7</sup>UK EQUATOR Centre, Centre for Statistics in Medicine, University of Oxford. Oxford. UK

<sup>8</sup>Jawaharlal Institute of Postgraduate Medical Education and Research, Puducherry, India

<sup>9</sup>Office of Science Dissemination, Centers for Disease Control and Prevention, Atlanta, GA, USA

<sup>10</sup>Department of Biostatistics and Epidemiology, School of Public Health, Center for Pharmacoepidemiology and Treatment Science, Rutgers University, New Brunswick, NJ, USA

11 JAMA Network Open, Chicago, IL, USA

 $^{12}\mathrm{Centre}$  for Health Research and Development, Society for Applied Studies, New Delhi, India

 $^{13}$ Child Health Evaluation Services, The Hospital for Sick Children Research Institute, Toronto, ON, Canada

 $^{14}\mbox{Department}$  of Psychiatry, University of Toronto, Toronto, ON, Canada

 $^{\rm 15}{\rm Aberdeen}$  Centre for Evaluation, University of Aberdeen, Aberdeen, UK

<sup>16</sup>Project PINK BLUE - Health & Psychological Trust Centre, Utako, Abuja, Nigeria

 $^{\rm 17}{\rm Department}$  of Sociology and Gerontology, Miami University, OH, USA

 $^{18}\mbox{Department}$  of Medical Statistics, London School of Hygiene and Tropical Medicine, London, UK

<sup>19</sup>Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford, UK

<sup>20</sup>Ottawa Hospital Research Institute, Ottawa, ON, Canada

<sup>21</sup>Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

 $^{22}\mbox{Department}$  of Epidemiology and Population Health, Stanford University, Palo Alto, CA, USA

<sup>23</sup>Institute for Evidence-Based Healthcare, Faculty of Health Sciences and Medicine, Bond University, University Drive, Robina, QLD, Australia

<sup>24</sup>Departments of Medicine, of Epidemiology and Population Health, of Biomedical Data Science, and of Statistics, and Meta-Research Innovation Center at Stanford (METRICS), Stanford University, Stanford. CA. USA

<sup>25</sup>MRC Clinical Trials Unit at University College London, London, UK
<sup>26</sup>University College London, UCL Great Ormond Street Institute of Child Health, London, UK

<sup>27</sup>NIHR Exeter Biomedical Research Centre, Faculty of Health and Life Sciences, University of Exeter, Exeter, UK

<sup>28</sup>Edinburgh Clinical Trials Unit, Usher Institute-University of Edinburgh, Edinburgh BioQuarter, Edinburgh, UK

<sup>29</sup>The BMJ, BMA House, London, UK

<sup>30</sup>Harvard Medical School, Boston, MA, USA

<sup>31</sup>Université Paris Cité, Inserm, INRAE, Centre de Recherche Epidémiologie et Statistiques, Université Paris Cité, Paris, France

<sup>32</sup>Clinical Trials Ontario, MaRS Centre, Toronto, ON, Canada

<sup>33</sup>Duke Clinical Research Institute, Duke University Medical Center, Durham, NC, USA

 $^{34}\mbox{Department}$  of Emergency Medicine, University of California, Los Angeles, CA, USA

<sup>35</sup>South African Medical Research Council, Cape Town, South Africa
<sup>36</sup>Warwick Applied Health, Warwick Medical School, University of Warwick, Coventry, UK

<sup>37</sup>MRC/CSO Social and Public Health Sciences Unit & Robertson Centre for Biostatistics, Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK

<sup>38</sup>Department of Health Research Methods Evidence and Impact, McMaster University, Hamilton, ON, Canada

<sup>39</sup>St Joseph's Healthcare Hamilton, Hamilton, ON, Canada

<sup>40</sup>York Trials Unit, Department of Health Sciences, University of York, York, UK

<sup>41</sup>Faculty of Medicine and Dentistry, University of Alberta, Edmonton, AB, Canada

<sup>42</sup>Centre for Evidence-Based Medicine Odense and Cochrane Denmark, Department of Clinical Research, University of Southern Denmark, Odense, Denmark

<sup>43</sup>Open Patient data Explorative Network, Odense University Hospital, Odense, Denmark

SPIRIT-CONSORT executive group: Isabelle Boutron, Université Paris Cité, France; An-Wen Chan, University of Toronto, Canada; Sally Hopewell, University of Oxford, UK; Asbjørn Hróbjartsson, University of Southern Denmark, Odense, Denmark; David Moher, University of Ottawa, Canada; Kenneth Schulz, University of North Carolina at Chapel Hill, USA. Gary Collins and Ruth Tunn, both of University of Oxford, UK, were also involved in leading the SPIRIT and CONSORT 2025 update.

SPIRIT-CONSORT 2025 consensus meeting participants: Rakesh Aggarwal, Jawaharlal Institute of Postgraduate Medical Education and Research, India; Michael Berkwits, Office of the Centers for Disease Control and Prevention, USA (formally JAMA and the JAMA Network at time of consensus meeting); Jesse A Berlin, Rutgers University/ JAMA Network Open USA; Nita Bhandari, Society for Applied Studies, India; Nancy J Butcher, The Hospital for Sick Children, Canada; Marion K Campbell, University of Aberdeen, UK; Runcie CW Chidebe, Project PINK BLUE, Nigeria/Miami University, Ohio, USA; Diana Elbourne, London School of Hygiene and Tropical Medicine, UK; Andrew J Farmer, University of Oxford, UK; Dean A Fergusson, Ottawa Hospital Research Institute, Canada; Robert M Golub, Northwestern University, USA; Steven N Goodman, Stanford University, USA; Tammy C Hoffmann, Bond University, Australia; John PA Ioannidis, Stanford University, USA; Brennan C Kahan, University College London, UK; Rachel L Knowles, University College London, UK: Sarah E Lamb. University of Exeter, UK; Steff Lewis, University of Edinburgh, UK; Elizabeth Loder, The BMJ, UK; Martin Offringa, Hospital for Sick Children Research Institute, Canada; Dawn P Richards, Clinical Trials Ontario, Canada: Frank W Rockhold, Duke University, USA: David L Schriger, University of California, USA; Nandi L Siegfried, South African Medical Research Council, South Africa; Sophie Staniszewska, University of Warwick, UK; Rod S Taylor, University of Glasgow, UK; Lehana Thabane, McMaster University/St Joseph's Healthcare, Canada; David Torgerson, University of York, UK; Sunita Vohra. University of Alberta, Canada; and Ian R White, University College

We dedicate SPIRIT 2025 to the late Doug Altman who was instrumental in the development of the SPIRIT and CONSORT statements and whose files and correspondence contributed to this update following his death. We gratefully acknowledge the contributions of all those who participated in the SPIRIT-CONSORT 2025 update Delphi survey. We also acknowledge Camilla Hansen Nejstgaard for conducting the scoping review to identify suggested changes to SPIRIT 2013 and for comparing SPIRIT 2025 with RoB2, Lasse Østengaard for developing the SCEB (SPIRIT-CONSORT Evidence Bibliographic) database of empirical evidence to support the development of SPIRIT 2025 and for assisting with handling references, Lina Ghosn for supporting the drafting of the expanded checklist, and Jen de Bayer and Patricia Logullo for their involvement during the consensus meeting. The SPIRIT 2025 statement is being simultaneously published in The BMJ, JAMA, The Lancet, Nature Medicine, and PLOS Medicine.

Contributors: A-WC, IB, SH, AH, DM, and KFS were responsible for the concept; SH, IB, A-WC, GSC, AH, KFS, and DM were responsible for the funding. A-WC, IB, SH, AH, GSC, DM, and KFS drafted specific sections of the manuscript. A-WC was guarantor of the work. All authors then critically revised the manuscript for important intellectual content and approved the final version. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

**Funding:** The 2025 update of SPIRIT and CONSORT was funded by an MRC-NIHR: Better Methods, Better Research grant (MR/W020483/1). The funder reviewed the design of the study but had no role in the collection, analysis or interpretation of the data; in the writing of the report; or in the decision to submit the article for publication. The views expressed are those of the authors and not necessarily those of the NIHR, the MRC, or the Department of Health and Social Care.

Competing interests: All authors have completed the ICMJE uniform disclosure form at www.icmje.org/disclosure-of-interest/ and declare: support from MRC-NIHR for the submitted work; SH, IB, A-WC, AH, KFS, and DM are members of the SPIRIT-CONSORT executive group. SH, IB, A-WC, AH, KFS, GSC, DM, MKC, NJB, MO, RST, and SV are involved in the development, update, implementation, and dissemination of several reporting guidelines. GSC is the director

of the LIK FOLIATOR Centre and a statistical editor for The BMI DM is the director of the Canadian EQUATOR Centre, and member of The BMI's regional advisory board for North America, IB is deputy director and PR is director of the French EQUATOR Centre, TCH is director of the Australasian EQUATOR Centre, JPAI is director of the US EQUATOR Centre, RA is president of the World Association of Medical Editors. MKC is chair of the MRC-NIHR: Better Methods Better Research funding panel. RCWC is executive director of Project PINK-BLUE which receives funding from Roche-Product. AF is director of the UK National Institute for Health and Care Research Health Technology Assessment Programme. DPR is a full time employee of FiveO2 Laboratories, which under contract to Clinical Trials Ontario provides services related to patient and public engagement; and is the volunteer vice president of the Canadian Arthritis Patient Alliance, which receives funding through independent grants from pharmaceutical companies. DLS is JAMA Associate Editor and receives editing stipends from JAMA and Annals of Emergency Medicine. IRW was supported by the MRC Programmes MCUU00004/07 and MCUU00004/09.

Ethical approval: Ethics approval was granted by the Central University Research Ethics Committee, University of Oxford (R76421/RE001). All Delphi participants provided informed consent to participate.

Patient and public involvement: The SPIRIT 2025 checklist items and the explanations here were developed using input from an international Delphi survey and consensus meeting. The Delphi survey was advertised via established patient and public involvement (PPI) networks, and 17 respondents self-identified as a "patient or public representative" and completed the Delphi survey. In addition, three participants in the expert consensus meeting were patient or public representatives who were leaders in advancing PPI.

Dissemination to participants and related patient and public communities: SPIRIT 2025 will be disseminated via a new website, consort-spirit.org, which will include materials designed for patients and the public.

**Provenance and peer review:** Not commissioned; externally peer reviewed.

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: http://creativecommons.org/licenses/by/4.0/.

- 1 Altman DG. Better reporting of randomised controlled trials: the CONSORT statement. BMJ 1996;313:570-1. doi:10.1136/ bmj.313.7057.570
- 2 Strengthening the credibility of clinical research. Lancet 2010;375:1225. doi:10.1016/S0140-6736(10)60523-5
- 3 Jones G, Abbasi K. Trial protocols at the BMJ. BMJ 2004;329:1360. doi:10.1136/bmi.329.7479.1360
- 4 Li T, Boutron I, Al-Shahi Salman R, et al. Review and publication of protocol submissions to Trials - what have we learned in 10 years? *Trials* 2016;18:34. doi:10.1186/s13063-016-1743-0
- 5 Krleza-Jerić K, Chan A-W, Dickersin K, Sim I, Grimshaw J, Gluud C. Principles for international registration of protocol information and results from human trials of health related interventions: Ottawa statement (part 1). BMJ 2005;330:956-8. doi:10.1136/bmj.330.7497.956
- 6 Lassere M, Johnson K. The power of the protocol. *Lancet* 2002; 360:1620-2. doi:10.1016/S0140-6736(02)11652-7
- 7 Chan A-W, Hróbjartsson A. Promoting public access to clinical trial protocols: challenges and recommendations. *Trials* 2018;19:116. doi:10.1186/s13063-018-2510-1
- 8 Zarin DA, Fain KM, Dobbins HD, Tse T, Williams RJ. 10-year update on study results submitted to ClinicalTrials.gov. N Engl J Med 2019;381:1966-74. doi:10.1056/NEJMsr1907644
- 9 Public Citizen Health Research Group v. Food and Drug Administration, 964 F Supp. 413 (DDC 1997).
- 10 Gryaznov D, von Niederhäusern B, Speich B, et al. Reporting quality of clinical trial protocols: a repeated cross-sectional study about the Adherence to SPIrit Recommendations in Switzerland, CAnada and GErmany (ASPIRE-SCAGE). BMJ Open 2022;12:e053417. doi:10.1136/bmjopen-2021-053417
- 11 Speich B, Odutayo A, Peckham N, et al. A longitudinal assessment of trial protocols approved by research ethics committees: The Adherance to SPIrit REcommendations in the UK (ASPIRE-UK) study. *Trials* 2022;23:601. doi:10.1186/s13063-022-06516-1
- 12 Kasenda B, von Elm E, You JJ, et al. Agreements between industry and academia on publication rights: a retrospective study of protocols and publications of randomized clinical trials. *PLoS Med* 2016;13:e1002046. doi:10.1371/journal. pmed.1002046

- 13 Getz K, Smith Z, Botto E, Murphy E, Dauchy A. New benchmarks on protocol amendment practices, trends and their impact on clinical trial performance. *Ther Innov Regul Sci* 2024;58:539-48. doi:10.1007/s43441-024-00622-9
- 14 Chan A-W, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013;158:200-7. doi:10.7326/0003-4819-158-3-201302050-00583
- 15 Chan A-W, Tetzlaff JM, Gøtzsche PC, et al. SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. BMJ 2013;346:e7586. doi:10.1136/bmj.e7586
- 16 Hopewell S, Chan A-W, Collins GS, et al. CONSORT 2025 statement: updated guideline for reporting randomised trials. BMJ 2025;389:e081123. doi:10.1136/bmj-2024-081123.
- 17 United Nations Educational Scientific Cultural Organization. UNESCO Recommendation on Open Science, 2021, Accessed at https:// unesdoc.unesco.org/ark:/48223/pf0000379949 on September 09, 2024
- Hopewell S, Boutron I, Chan A-W, et al. An update to SPIRIT and CONSORT reporting guidelines to enhance transparency in randomized trials. *Nat Med* 2022;28:1740-3. doi:10.1038/s41591-022-01989-8
- 19 Hopewell S, Chan A-W, Boutron I, et al. Protocol for updating the SPIRIT 2013 (Standard Protocol Items: Recommendations for Interventional Trials) and CONSORT 2010 (CONsolidated Standards Of Reporting Trials) Statements (version 1.0, August 2022). OSF. IO/6HJYG.
- 20 Tunn R, Boutron I, Chan A-W, et al. Methods used to develop the SPIRIT 2024 and CONSORT 2024 Statements. J Clin Epidemiol 2024;169:111309. doi:10.1016/j.jclinepi.2024.111309
- 21 Moher D, Schulz KF, Simera I, Altman DG. Guidance for developers of health research reporting guidelines. *PLoS Med* 2010;7:e1000217. doi:10.1371/journal.pmed.1000217
- 22 Nejstgaard CH, Boutron I, Chan A-W, et al. A scoping review identifies multiple comments suggesting modifications to SPIRIT 2013 and CONSORT 2010. *J Clin Epidemiol* 2023;155:48-63. doi:10.1016/j.jclinepi.2023.01.003
- 23 Østengaard L, Barrientos A, Boutron I, et al. Development of a topic-specific bibliographic database supporting the updates of SPIRIT 2013 and CONSORT 2010. Cochrane Evidence Synthesis and Methods. 2024;2:e12057. doi:10.1002/cesm.12057
- 24 Junqueira DR, Zorzela L, Golder S, et al, CONSORT Harms Group. CONSORT Harms 2022 statement, explanation, and elaboration: updated guideline for the reporting of harms in randomized trials. J Clin Epidemiol 2023;158:149-65. doi:10.1016/j. iclinepi.2023.04.005
- 25 Butcher NJ, Monsour A, Mew EJ, et al. Guidelines for reporting outcomes in trial protocols: The SPIRIT-Outcomes 2022 Extension. JAMA 2022;328:2345-56. doi:10.1001/jama.2022.21243
- 26 Boutron I, Altman DG, Moher D, Schulz KF, Ravaud P, CONSORT NPT Group. CONSORT Statement for randomized trials of nonpharmacologic treatments: A 2017 update and a CONSORT extension for nonpharmacologic trial abstracts. *Ann Intern* Med 2017:167:40-7. doi:10.7326/M17-0046
- 27 Hoffmann TC, Glasziou PP, Boutron I, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;348:g1687. doi:10.1136/ bmj.g1687
- 28 Hróbjartsson A, Boutron I, Collins GS, et al. SPIRIT 2025 Explanation and Elaboration: updated guideline for protocols of randomised trials. BMJ 2025;389:e081660. doi:10.1136/bmj-2024-081660
- 29 Barnes C, Boutron I, Giraudeau B, Porcher R, Altman DG, Ravaud P. Impact of an online writing aid tool for writing a randomized trial report: the COBWEB (Consort-based WEB tool) randomized controlled trial. BMC Med 2015;13:221. doi:10.1186/s12916-015-0460-y
- 30 Chauvin A, Ravaud P, Moher D, et al. Accuracy in detecting inadequate research reporting by early career peer reviewers using an online CONSORT-based peer-review tool (COBPeer) versus the usual peer-review process: a cross-sectional diagnostic study. BMC Med 2019;17:205. doi:10.1186/s12916-019-1436-0
- 31 Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. Syst Rev 2021;10:89. doi:10.1186/s13643-021-01626-4
- World Medical Association. World Medical Association Declaration of Helsinki: Ethical Principles for Medical Research Involving Human Participants. JAMA 2025;333:71-4. doi:10.1001/ jama.2024.21972
- 33 Getz KA, Stergiopoulos S, Short M, et al. The impact of protocol amendments on clinical trial performance and cost. *Ther Innov Regul* Sci 2016;50:436-41. doi:10.1177/2168479016632271
- 34 Orkin AM, Gill PJ, Ghersi D, et al, CONSERVE Group. Guidelines for reporting trial protocols and completed trials modified due to the COVID-19 pandemic and other extenuating circumstances: The CONSERVE 2021 Statement. JAMA 2021;326:257-65. doi:10.1001/ jama.2021.9941

- 35 Chan AW, Karam G, Pymento J, et al. Reporting summary results in clinical trial registries: updated guidance from WHO. Lancet Glob Health 2025;13:e759-68. doi:10.1016/S2214-109X(24)00514-X.
- 36 International Council for Harmonisation of Technical Requirements for Pharmaceuticals for Human Use. International Council for Harmonisation (ICH) harmonised guideline: Guideline for good clinical practice, E6(R3). 6 January 2025. https://www.ich.org/page/ efficacy-guidelines
- 37 Gamble C, Krishan A, Stocken D, et al. Guidelines for the content of statistical analysis plans in clinical trials. JAMA 2017;318:2337-43. doi:10.1001/jama.2017.18556
- 38 World Health Organization. Guidance for best practices for clinical trials. 2024. Accessed at https://www.who.int/publications/i/ item/9789240097711 on September 09, 2024
- 39 Chan A-W, Pello A, Kitchen J, et al. Association of trial registration with reporting of primary outcomes in protocols and publications. JAMA 2017;318:1709-11. doi:10.1001/jama.2017.13001
- 40 Kahan BC, Hall SS, Beller EM, et al. Consensus statement for protocols of factorial randomized trials: extension of the SPIRIT 2013 Statement. JAMA Netw Open 2023;6:e2346121. doi:10.1001/ jamanetworkopen.2023.46121
- 41 Calvert M, Kyte D, Mercieca-Bebber R, et al, the SPIRIT-PRO Group. Guidelines for inclusion of patient-reported outcomes in clinical trial protocols: The SPIRIT-PRO Extension. *JAMA* 2018;319:483-94. doi:10.1001/jama.2017.21903
- 42 Porcino AJ, Shamseer L, Chan A-W, et al, SPENT group. SPIRIT extension and elaboration for n-of-1 trials: SPENT 2019 checklist. BMJ 2020;368:m122. doi:10.1136/bmj.m122
- 43 Yap C, Rekowski J, Ursino M, et al. Enhancing quality and impact of early phase dose-finding clinical trial protocols: SPIRIT Dose-finding Extension (SPIRIT-DEFINE) guidance. *BMJ* 2023;383:e076386. doi:10.1136/bmj-2023-076386
- 44 Kendall TJ, Robinson M, Brierley DJ, et al, SPIRIT-Path group. Guidelines for cellular and molecular pathology content in clinical trial protocols: the SPIRIT-Path extension. *Lancet* Oncol 2021;22:e435-45. doi:10.1016/S1470-2045(21)00344-2
- 45 Cruz Rivera S, Liu X, Chan A-W, Denniston AK, Calvert MJ, SPIRIT-AI and CONSORT-AI Working GroupSPIRIT-AI and CONSORT-AI Steering GroupSPIRIT-AI and CONSORT-AI Consensus Group. Guidelines for clinical trial protocols for interventions involving artificial intelligence: the SPIRIT-AI extension. *Nat Med* 2020;26:1351-63. doi:10.1038/s41591-020-1037-7

- 46 Dai L, Cheng CW, Tian R, et al. Standard Protocol Items for Clinical Trials with Traditional Chinese Medicine 2018: Recommendations, Explanation and Elaboration (SPIRIT-TCM Extension 2018). Chin J Integr Med 2019;25:71-9. doi:10.1007/s11655-018-2999-x
- 47 Manyara AM, Davies P, Stewart D, et al. Reporting of surrogate endpoints in randomised controlled trial protocols (SPIRIT-Surrogate): extension checklist with explanation and elaboration. BMJ 2024;386:e078525. doi:10.1136/bmj-2023-078525
- 48 Blanco D, Donadio MVF, Cadellans-Arróniz A. Enhancing reporting through structure: a before and after study on the effectiveness of SPIRIT-based templates to improve the completeness of reporting of randomized controlled trial protocols. Res Integr Peer Rev 2024;9:6. doi:10.1186/s41073-024-00147-7
- 49 Tan ZW, Tan AC, Li T, et al. Has the reporting quality of published randomised controlled trial protocols improved since the SPIRIT statement? A methodological study. *BMJ Open* 2020;10:e038283. doi:10.1136/bmjopen-2020-038283
- 50 Happo SM, Halkoaho A, Lehto SM, Keränen T. The effect of study type on research ethics committees' queries in medical studies. Res Ethics 2017;13:115-27. doi:10.1177/1747016116656912
- 51 Russ H, Busta S, Jost B, Bethke TD. Evaluation of clinical trials by Ethics Committees in Germany--results and a comparison of two surveys performed among members of the German Association of Research-Based Pharmaceutical Companies (vfa). Ger Med Sci 2015;13:Doc02.
- 52 TARG Meta-Research Group & Collaborators. Estimating the prevalence of discrepancies between study registrations and publications: a systematic review and meta-analyses. *BMJ Open* 2023;13:e076264. doi:10.1136/bmjopen-2023-076264
- 53 Chan A-W, Hróbjartsson A, Haahr MT, Gøtzsche PC, Altman DG. Empirical evidence for selective reporting of outcomes in randomized trials: comparison of protocols to published articles. JAMA 2004;291:2457-65. doi:10.1001/jama.291.20.2457
- 54 Schönenberger CM, Griessbach A, Taji Heravi A, et al. A metaresearch study of randomized controlled trials found infrequent and delayed availability of protocols. *J Clin Epidemiol* 2022;149:45-52. doi:10.1016/j.jclinepi.2022.05.014
- 55 Spence O, Hong K, Onwuchekwa Uba R, Doshi P. Availability of study protocols for randomized trials published in high-impact medical journals: A cross-sectional analysis. Clin Trials 2020;17:99-105. doi:10.1177/1740774519868310

Web appendix: Supplementary materials