



CIRCULAR

Imphal, the 18th March, 2026

No. 55/RIMS-MRU/2026: It is to notify that, the 10th **Research Masterclasses 2026**, of the Department of Health Research, Ministry of Health and Family Welfare, Government of India, will be conducted virtually, on **27th March, 2026 (Friday)**.

2. All the faculties (RIMS, Dental College, and College of Nursing), members of EC, LRAC of MRU, Principal Investigators undertaking MRU funding projects (including under process projects) and residents are invited to attend the session at **Sushruta Hall, RIMS, Imphal**.

Date: 27.03.2026 (Friday)


Time: 3:00 PM onwards

Venue: Sushruta Hall, RIMS, Imphal (**Designated Site for Participation for RIMS**)

Event name: Research Masterclass under DHR-ICMR Research Grand Rounds

Speaker Name: Dr. Manoj V. Murhekar, Director & Scientist 'G', National Institute of Epidemiology, Chennai, Tamil Nadu.

3. The **research paper** to be discussed during the Masterclass will be uploaded on the **RIMS website** and circulated to the concerned Departments/Colleges through official email.


(Prof. Ng. Gunindro Singh)

Nodal Officer (i/c)

Multi-Disciplinary Research Unit,
RIMS, Imphal

Copy to:

1. The P.S. to Director, RIMS, for kind information of Director
2. The P.A. to Medical Superintendent, RIMSH, for kind information
3. The Dean (Academic), RIMS, for kind information.
4. The Principal, Dental College, RIMS
5. The Principal, College of Nursing, RIMS
6. The Head of Department, RIMS, Imphal
7. The Chairperson/Co-Chairperson/Member, LRAC, MRU, RIMS.....
8. The Member, EC, MRU, RIMS, Imphal.....
9. The Principal Investigator, RIMS
- ✓ 10. The IT Cell, RIMS – with a request for uploading the notice in the website & technical support on **27.03.26**
11. Asst. Engineer (Elect. /Civil), RIMS - with a request for ensuring uninterrupted power supply & optimum AC functioning.
12. The Care Taker, Sushruta Hall, RIMS, Imphal- for proper upkeep of the venue & accompanying facilities.
13. Guard file.

No. R.11016/03/2025-HR
भारत सरकार/Government of India
स्वास्थ्य एवं परिवार कल्याण मंत्रालय/Ministry of Health & Family Welfare
स्वास्थ्य अनुसंधान विभाग/Department of Health Research

2nd Floor, IRCS Building
Sansad Marg, New Delhi – 110001
Dated 12.03.2026

To
The Dean/ Principal/ Director of Medical Colleges/ Institutes

Subject: Request to attend Research Masterclasses for MRU network– reg.

Sir/Madam,

DHR-ICMR has initiated a dedicated platform to conduct Research Grand Rounds to strengthen the National research ecosystem through sustained collaboration and knowledge exchange. The objectives of the Research Grand Rounds are as follows:

- I. To deliberate on research methodologies, analytical tools, and emerging scientific approaches
- II. To strengthen the methodological understanding amongst researchers needed to implement different kinds of research.
- III. To foster collaboration and connectivity across research institutions

2. These Research Grand Rounds will be organized as monthly webinars entitled 'Research Masterclass' proposed around the last Friday of each month. The speakers for these Research Masterclasses will be eminent research scientists in the country who will be discussing their original research work in details from methodological point of view.

3. The next Research Masterclass is scheduled for **27.03.2026 (Friday) at 3:00 PM**. The invited speaker is **Dr. Manoj V. Murhekar, Director & Scientist 'G'**, National Institute of Epidemiology, Chennai, Tamil Nadu. The research paper to be discussed during the research masterclass is enclosed. The link for the research masterclass will be shared shortly.

4. Accordingly, it is requested to kindly disseminate the information in your institution and ensure maximum participation in Research Masterclass. Your institute is requested to share at least two questions related to research paper attached on the following email: **dhr-mru@gov.in** latest by 24.03.2026. These questions will be discussed with the speaker during masterclass.

Yours faithfully,



(Dharkat R. Luikang)

Deputy Secretary to the Govt. of India

Copy to: The Nodal Officer of Multi-Disciplinary Research Units (MRUs)

THE LANCET

Infectious Diseases

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Thangaraj JWV, Krishna NS, Devika S, et al. Estimates of the burden of human rabies deaths and animal bites in India, 2022–23: a community-based cross-sectional survey and probability decision-tree modelling study. *Lancet Infect Dis* 2024; published online Sept 30. [https://doi.org/10.1016/S1473-3099\(24\)00490-0](https://doi.org/10.1016/S1473-3099(24)00490-0).

SUPPLEMENTARY APPENDIX

SUPPLEMENT TO:


Human rabies death and animal bite burden estimates in India, 2022-2023

Appendix

Contents No.	Page
1. Figure 1: Districts selected for the Human rabies death and animal bite burden survey, India, 2022-2023.....	1
2. Weight calculation, Human rabies death and animal bite burden survey, India, 2022-2023	2
3: Steps and sources of input data for the WHO decision tree probability model, Human rabies death and animal bite burden survey, India, 2022-2023	3
4. Figure 2: Flow chart illustrating the enrolment of participants in the survey.....	5
5. Table 1: Characteristics of Animal Bite among study participants, Human rabies death and animal bite burden survey, India, 2022-2023	6
6. Figure 3: Annual Incidence of Dog bite by Regions, Human rabies death and animal bite burden survey, India, 2022-2023	8
7. Table 2: Factors associated with non-receipt of ARV among dog bite victims, Human rabies death and animal bite burden survey, India, 2022-2023	9
8. Table 3: Household pet ownership and rabies vaccination, Human rabies death and animal bite burden survey, India, 2022-2023.....	12
9. Figure 4: Classification of bites as bite by a suspected rabid dog or normal dog using Tepsumethanon et al. Six criteria for rabies diagnosis in living dogs.....	13
10. Table 4: Dog Rabies positivity among suspected rabid dogs across laboratories of India (2017-2022)	14
11. Table 5: Estimates of deaths due to human rabies using the WHO decision tree probability model (sensitivity analysis)	15
12. Table 6: COVID-19 movement restrictions overlapping the Recall period for animal bite history one year, Human rabies death and animal bite burden survey, India, 2022-2023.....	16
13. Table 7a: Comparison of ARV vaccination by recall and documented evidence.....	17
14. Figure 5: Flow chart illustrating the enrolment of participants in the survey	18
15. Table 8: Socio-demographic characteristics of study participants by regions, Human rabies death and animal bite burden survey, India, 2022-2023	19
16. Table 9: Characteristics of Animal Bite among study participants by regions, Human rabies death and animal bite burden survey, India, 2022-2023	21
17. Table 10: Household pet ownership and rabies vaccination by regions, Human rabies death and animal bite burden survey, India, 2022-2023	30
18. Table 11: Comparison of the age, gender and type of residence of the participants in the study with that of the Census 2011 India population.....	32
19. STROBE Statement—Checklist of items that should be included in reports of cross-sectional studies	33

1. Figure 1: Districts selected for the Human rabies death and animal bite burden survey, India, 2022-2023

Region	State	Districts
North	Uttarakhand	Dehradun/Pithoragarh/Udham Singh Nagar/Haridwar
	Uttar Pradesh	Meerut/Sitapur/Allahabad/Mau
	Punjab	Gurdaspur/Shahid Bhagat Singh Nagar/Firozpur/Amritsar
North-East	Nagaland	Mon/Zunheboto/Dimapur/Kiphire
	Tripura	West Tripura/Sepahijala/Gomati/Unakoti
	Assam	Barpeta/Dhemaji/Cachar/Nalbari
East	Bihar	Madhubani/Siwan/Nalanda/Nawada
	Chhattisgarh	Baloda Baazar/Bastar/Korba/Rajnandgaon
	Odisha	Sundargarh/Jagatsinghapur/Khordha/Kalahandi
West	Madhya Pradesh	Bhind/Neemuch/Rajgarh/Chhindwara
	Gujarat	Mahesana/Rajkot/Kheda/Valsad
	Maharashtra	Nagpur/Nashik/Raigarh/Satara
South	Telangana	Kamareddy, Hyderabad, Narayanpet, Mahabubabad
	Kerala	Kannur/Malappuram/Ernakulam/Pathanamthitta
	Tamil Nadu	Chennai/Salem/Nagapattinam/Tirunelveli



2. Weight calculation, Human rabies death and animal bite burden survey, India, 2022-2023

We used multistage sampling comprising different stages of selection involving varying probabilities of selection. We have accounted for the different probabilities in terms of weights in the final analysis.

The steps involved in the calculation of weight:

1. Determining the probability of selection at each stage:

At each stage of sampling, we calculate the probability of selection for each unit (i.e., state → district → village/wards → CEB). The probability of selection at each stage (P_i) depends on the sampling method we used (i.e., simple random sampling in the selection of states and probability proportional to size in subsequent stages).

2. Calculation of overall probability of selection:

We calculated the overall probability of selection for each sampled unit as the product of the probabilities of selection at each stage as $P_{\text{overall}} = P_{\text{state}} \times P_{\text{district}} \times P_{\text{village/ward}} \times P_{\text{CEB}}$. Here, P_{state} is the probability of selecting a state (primary sampling unit), P_{district} is the probability of selecting a district (secondary sampling unit) within the state, and so on.

3. Computing Design Weights:

The design weight (W) for each unit is calculated as the inverse of its overall probability of selection i.e., $W = 1 / P_{\text{overall}}$

4. Adjust for Nonresponse:

Based on the response rates we calculated the nonresponse adjustment factor. We then adjust the weights of the respondents using a nonresponse adjustment factor to compensate for the non-response and calculated the final sampling weight.

3: Steps and sources of input data for the WHO decision tree probability model, Human rabies death and animal bite burden survey, India, 2022-2023

The model uses two formulas to predict the number of human deaths due to rabies.

The formula to calculate human rabies deaths is

Incidence of suspected rabies dog bite ($I_{\text{Rabid dog bite}}$) * Population at risk (Q) * Probability of death after being bitten by a rabid dog (P_{Death})

Step 1: Incidence of suspected rabies dog bite ($I_{\text{Rabid dog bite}}$)

Dog bites identified in the survey was classified as bite by a normal dog or bite by a suspected rabid dog using the Tepsumethanon et al Six criteria for rabies diagnosis in living dogs. Incidence of suspected rabies dog bite was calculated by considering the number of suspected rabid dog bites in last one year as the numerator and surveyed population as the denominator.

Step 2: Population at risk (Q)

Population at risk (Q) was the population of India obtained from the 2021 census report


Step 3: Probability of death after being bitten by a rabid dog (P_{Death})

Probability of death after being bitten by a rabid dog (P_{Death}) was calculated using the following formula from the decision tree model

$$P1 * [(P2 * P6) + (P3 * P7) + (P4 * P8) + P5 * P9] * (1 - P10)$$

Probabilities	Source of the input parameter/data
P1 Suspected rabid dog being rabid	Using data from the 6 animal laboratories for 12 states of India
P2 Bite injury to the head	Using the data from community survey
P3 Bite injury to the arms	Using the data from community survey
P4 Bite injury to the trunk	Using the data from community survey
P5 Bite injury to the legs	Using the data from community survey
P6 Development of rabies following bite injury to the head	
P7 Development of rabies following bite injury to the arms	Using the data from literature
P8 Development of rabies following bite injury to the trunk	
P9 Development of rabies following bite injury to the leg	
P10 Probability of receiving post-exposure treatment if bitten by a suspected rabid dog	Using the data from community survey

4. Figure 2: Flow chart illustrating the enrolment of participants in the survey




5. Table 1: Characteristics of Animal Bite among study participants, Human rabies death and animal bite burden survey, India, 2022-2023

Characteristics of animal and dog bite	Number
No of animal bites reported in the last 5 years	5,309
No of dog bites reported in the last 5 year	4,289
No of animal bites reported in the last 3 years	4,073
No of dog bites reported in the last 3 year	3,218
No of animal bites reported in the last 1 year	2,052
No of dog bites reported in the last 1 year	1,576
Annual incidence	Incidence (Per 1000) 95% CI
Unweighted annual incidence of animal bite	6.1 [5.8-6.3]
Weighted annual incidence of animal bite	6.6 [5.7-7.6]
Unweighted annual incidence of dog bite	4.7 [4.4-4.9]
Weighted annual incidence of dog bite	5.6 [4.8-6.6]
Weighted annual incidence of dog bite by age (in years) of the bite victim	
0-14	7.5 [5.8-9.6]
15-60	4.7 [4.1-5.4]
>60	7.8 [5.2-11.5]
Weighted annual incidence of dog bite by gender of the bite victim	
Male	7.6 [6.6-8.8]
Female	3.5 [2.8-4.4]
Weighted annual incidence of dog bite by type of residence	
Rural	5.8 [4.8-6.9]
Urban	5.0 [4.1-6.0]
Dog bite victims sought medical care at a health facility (n=1,576)	
Yes	1,317 (83.6)
Time gap between the bite and medical care sought (n=1,317)	
Within 1 day	1,067 (81.0)
1 day – 3 days	236 (17.9)
3 days – 7 days	8 (0.6)
More than 7 days	6 (0.5)
Anti Rabies Vaccine coverage of dog bite victims	
Received anti rabies vaccine (ARV) (n=1,576)	1,253 (79.5)

Characteristics of animal and dog bite	Number (% of total)
Route of administration (n=1,253)	
Intra dermal (ID)	836 (66.7)
Intra muscular (IM)	417 (33.3)
ARV dose completed (ID route of administration) (n=836)	
1 dose only	55 (6.6)
2 doses only	96 (11.5)
3 doses only	181 (21.6)
4 doses	504 (60.3)
ARV dose completed (IM route of administration) (n=417)	
1 dose only	29 (6.9)
2 doses only	30 (7.2)
3 doses only	137 (32.9)
4 doses only	87 (20.9)
5 doses	134 (32.1)
Reason for not vaccinated (n=323)	
Not aware of post exposure prophylaxis/Don't know where to get the vaccine	183 (56.7)
Generally, don't believe in vaccine/Vaccine will not be helpful in preventing the disease	30 (9.3)
Consulted a traditional healer/Siddha/Ayurveda practitioner	24 (7.4)
Minor scratch	19 (5.9)
Can't miss work	15 (4.6)
Only TT taken	13 (4.0)
Couldn't afford the vaccine	10 (3.1)
Bitten by a vaccinated dog	9 (2.7)
Hospital too far/No transportation	7 (2.2)
Fear of side effects, pain at the injection site and fever/ Heard from social media/neighbor/religious leader/mass media that vaccine is harmful	7 (2.2)
Vaccine not available	6 (1.9)
Received rabies immunoglobulin (RIG) (n=1,576)	81 (5.1)
Site of RIG (n=81)	
Into the wound	48 (59.4)
Into the wound & Deltoid	28 (35.4)
Into the wound & Gluteal	2 (2.5)
Into the wound & Antero lateral thigh	3 (3.7)

6. Figure 3: Annual Incidence of Dog bite by Regions, Human rabies death and animal bite burden survey, India, 2022-2023



7. Table 2: Factors associated with non-receipt of ARV among dog bite victims, Human rabies death and animal bite burden survey, India, 2022-2023

Characteristics	Vaccinated		OR (95% CI)	P Value	Adjusted OR (95% CI)	P Value
	No (n= 323 (%))	Yes (n=1,253 (%))				
Region						
North	95 (29.4)	314 (25.1)	4.31 [2.60 - 7.13]	<0.0001	5.25 [3.02 - 9.13]	<0.0001
North-East	87 (26.9)	174 (13.9)	7.12 [4.07 - 12.45]	<0.0001	5.42 [2.94 - 10.00]	<0.0001
East	33 (10.2)	194 (15.5)	2.42 [1.30 - 4.51]	0.005	2.42 [1.24 - 4.73]	0.010
West	87 (26.9)	272 (21.7)	4.55 [2.68 - 7.74]	<0.0001	4.66 [2.66 - 8.16]	<0.0001
South	21 (6.5)	299 (23.8)	1			
Age (in years)						
0-14	107 (33.1)	395 (31.5)	0.98 [0.63 - 1.53]	0.933		
15-60	179 (55.4)	724 (57.8)	0.89 [0.58 - 1.38]	0.617		
>60	37 (11.5)	134 (10.7)	1			
Gender						
Male	222 (68.7)	825 (65.8)	1.14 [0.87 - 1.50]	0.344		
Female	101 (31.3)	428 (34.2)	1			
Dog Type						
Pet	139 (43.0)	475 (37.9)	1.24 [0.94 - 1.63]	0.133	1.39 [1.02 - 1.89]	0.039
Stray/ Wild	184 (57.0)	778 (62.1)	1			


Characteristics	Vaccinated		OR (95% CI)	P Value	Adjusted OR (95% CI)	P Value
	No (n= 323 (%))	Yes (n=1,253 (%))				
Status of Dog after ten days						
Alive	232 (71.8)	834 (66.6)	1.28 [0.97 - 1.70]	0.085	1.12 [0.82 - 1.53]	0.482
Died/Killed within 10 days/Don't Know/Not available for observation	91 (28.2)	419 (33.4)	1			
Dog vaccinated						
Yes	48 (14.9)	182 (14.5)	1.03 [0.70 - 1.50]	0.890		
No/Don't Know	275 (85.1)	1,071 (85.5)	1			
Total number of bite wounds						
1	305 (94.4)	1,152 (91.9)	1.49 [0.84 - 2.62]	0.170	1.35 [0.76 - 2.42]	0.308
≥2	18 (5.6)	101 (8.1)	1			
Type of wound						
Abrasion	244 (75.5)	786 (62.7)	1.83 [1.35 - 2.49]	<0.0001	2.00 [1.42 - 2.82]	<0.0001
Laceration/ Puncture Wound/Avulsion	79 (24.5)	467 (37.3)	1			
Site of bite						
Head/Face	12 (3.7)	30 (2.4)	1			
Trunk	10 (3.1)	59 (4.7)	0.42 [0.16 - 1.11]	0.080	0.54 [0.20 - 1.48]	0.231
Arm/Forearm	87 (26.9)	302 (24.1)	0.72 [0.34 - 1.53]	0.394	0.94 [0.42 - 2.10]	0.873
Leg/Feet	214 (66.2)	862 (68.8)	0.62 [0.31 - 1.25]	0.182	0.85 [0.40 - 1.80]	0.663

Characteristics	Vaccinated		OR (95% CI)	P Value	Adjusted OR (95% CI)	P Value
	No (n= 323 (%))	Yes (n=1,253 (%))				
Education of head of the household						
Professional or honors/graduate	29 (9.0)	132 (10.5)	1			
Schooling (primary, middle and high school)	218 (67.5)	918 (73.3)	1.08 [0.69 - 1.70]	0.737	0.86 [0.54 - 1.40]	0.551
Illiterate/ read and write with no formal education	76 (23.5)	203 (16.2)	1.70 [1.03 - 2.82]	0.038	1.51 [0.88 - 2.62]	0.136
Type of house						
Pucca	144 (44.6)	758 (60.5)	1			
Semi-pucca	82 (25.4)	336 (26.8)	1.28 [0.94- 1.75]	0.114	1.37 [0.95- 1.96]	0.088
Kuccha	97 (30.0)	159 (12.7)	3.21 [2.32 - 4.45]	<0.0001	2.64 [1.77 - 3.94]	<0.0001
Toilet facilities						
Yes (Own, Shared & Public toilet)	283 (87.6)	1,130 (90.2)	1			
No (Use open space)	40 (12.4)	123 (9.8)	1.30 [0.83 - 2.03]	0.250		
Cluster type						
Rural	246 (76.2)	868 (69.3)	1.42 [0.98 - 2.04]	0.062	1.07 [0.73 - 1.56]	0.716
Urban	77 (23.8)	385 (30.7)	1			

8. Table 3: Household pet ownership and rabies vaccination, Human rabies death and animal bite burden survey, India, 2022-2023

Characteristics of the pet/domestic animal	Number (% of total)
Owns pet/domestic animal (N=78,807)	19,994 (25.4)
Type of pet (N=19,994)	
Dog	3,714 (18.6)
Cat	2,817 (14.1)
Cow	10,231 (51.2)
Goat/sheep	5,137 (25.7)
Pig	1,637 (8.2)
Horse	31 (0.2)
Donkey	10 (0.1)
Rabbit	131 (0.7)
Monkey	8 (0.04)
Others (Buffalo, Ox, Camel, Mongoose)	2,966 (14.8)
No of pet dogs in each household (N=3,714)	
One dog	3,186 (85.8)
Two dogs	401 (10.8)
Three or more dogs	127 (3.4)
Vaccination status of the dog (N=4,369)	
Vaccinated	2,223 (50.9)
Not vaccinated	2,052 (47.0)
Don't know about vaccination status	94 (2.1)
Own dog(s) died in the past year (N=78,807)	
Yes	1,150 (1.5)
Provide care for other dogs (N=78,807)	
Stray dog	13,215 (16.8)
Community dog	7,217 (9.1)
Does not provide care	58,375 (74.1)
Type of care provide for the stray/community dogs (N=20,432)	
Food	19,759 (96.7)
Water	14,333 (70.2)
Shelter	953 (4.7)
Veterinary care	17 (0.1)
Any member in the household received pre-exposure prophylaxis (N=78,807)	
Received	453 (0.6)

9. Figure 4: Classification of bites as bite by a suspected rabid dog or normal dog using Tepsu Methanon et al. Six criteria for rabies diagnosis in living dogs



10. Table 4: Dog Rabies positivity among suspected rabid dogs across laboratories of India (2017-2022)

Name of the State	Year	No. of Positive samples	No. of Negative samples	Total no. of samples tested	Positivity	Region as per study
Karnataka	2017 - 2022	931	271	1,202	0.775	South Zone
Assam	2017 - 2022	15	2	17	0.882	North East Zone
Manipur	2017 - 2022	43	14	57	0.754	North East Zone
Himachal Pradesh	2018 & 2022	2	1	3	0.667	North Zone
Andhra Pradesh	2020	0	2	2	0.000	South Zone
Haryana	2020	0	1	1	0.000	North Zone
Gujarat	2017 - 2022	27	4	31	0.871	West Zone
Telangana	2022	1	0	1	1.000	South Zone
Tamil Nadu	2018 - 2021	90	36	126	0.714	South Zone
Maharashtra	2017 - 2022	85	46	131	0.649	West Zone
Kerala	2020 - 2022	200	61	261	0.766	South Zone
Uttar Pradesh	2017 - 2022	13	35	48	0.271	North Zone
		1,407	473	1,880	0.748	

11. Table 5: Estimates of deaths due to human rabies using the WHO decision tree probability model (sensitivity analysis)

Parameter	Description	Three-year survey data	One-year data Using Lower level 95% CI of P1	One year data Using Upper level 95% CI of P1
	No. of suspected rabid dog bites	164	69	69
P1	Probability of a suspected rabid dog being confirmed rabid on laboratory diagnosis (Binomial distribution: n=1,880)	0.748	0.728	0.768
P2	Bite injury to the head or neck (Point estimate)	0.0305	0.0290	0.0290
P3	Bite injury to the hand or arm (Point estimate)	0.3049	0.2754	0.2754
P4	Bite injury to the trunk (Point estimate)	0.0549	0.087	0.087
P5	Bite injury to the leg or foot (Point estimate)	0.6098	0.6087	0.6087
P6	Probability of developing rabies following a bite injury to the head by a rabid dog	Triangular distribution: minimum=0.30, mode=0.45, maximum=0.60		
P7	Probability of developing rabies following a bite injury to the hand or arm by a rabid dog	Triangular distribution: minimum=0.15, mode=0.28, maximum=0.40		
P8	Probability of developing rabies following a bite injury to the trunk by a rabid dog	Triangular distribution: minimum=0.00, mode=0.05, maximum=0.10		
P9	Probability of developing rabies following a bite injury to the leg or foot by a rabid dog	Triangular distribution: minimum=0.00, mode=0.05, maximum=0.10		
P10	Probability of an individual receiving anti-rabies vaccination if bitten by a suspected rabid dog (Point estimate)	0.8232	0.7826	0.7826
	Probability of death (P_{Death})	0.01750043	0.01977501	0.02086155
	Incidence of suspected rabid dog bite (Annual Incidence)	(164/337,808)/3	69/337,808	69/337,808
	At risk population (Q)	1,379,750,000	1,379,750,000	1,379,750,000
	No. of human rabies deaths (95% Uncertainty Limit) in a year	3,908 (2,741 - 5,016)	5,573 (3,897 - 7,165)	5,879 (4,078 - 7,560)

12. Table 6: COVID-19 movement restrictions overlapping the Recall period for animal bite history one-year, Human rabies death and animal bite burden survey, India, 2022-2023

State Name	Start date of the Survey	End date of the Survey	Recall period for animal bite history one year		Recall period overlapping with the COVID-19 movement restriction corresponding to first and second wave (March 2020 - August 2021)
Uttarakhand	19-10-2022	10-02-2023	19-10-2021	10-02-2022	No
Uttar Pradesh	24-02-2023	03-05-2023	24-02-2022	03-05-2022	No
Nagaland	23-06-2023	16-08-2023	23-06-2022	16-08-2022	No
Tripura	08-12-2022	04-04-2023	08-12-2021	04-04-2022	No
Assam	29-03-2023	26-08-2023	29-03-2022	26-08-2022	No
Bihar	22-12-2022	26-04-2023	22-12-2021	26-04-2022	No
Chhattisgarh	17-07-2023	25-08-2023	17-07-2022	25-08-2022	No
Odisha	24-11-2022	21-02-2023	24-11-2021	21-02-2022	No
Madhya Pradesh	08-02-2023	25-05-2023	08-02-2022	25-05-2022	No
Gujarat	23-11-2022	23-02-2023	23-11-2021	23-02-2022	No
Maharashtra	31-10-2022	19-01-2023	31-10-2021	19-01-2022	No
Telangana	05-01-2023	30-06-2023	05-01-2022	30-06-2022	No
Kerala	26-04-2023	29-06-2023	26-04-2022	29-06-2022	No
Tamil Nadu	02-03-2022	26-03-2022	02-03-2021	26-03-2021	Yes
Punjab	19-09-2022	10-02-2023	19-09-2021	10-02-2022	No

13. Table 7a: Comparison of ARV vaccination by recall and documented evidence

Arv dose completed	Provides document	Information based on recall
	(n =240)	(n=880)
	n %	n %
1 dose only	26 (10.8)	46 (5.2)
2 doses only	21 (8.7)	89 (10.1)
3 doses only	42 (17.5)	237 (26.9)
4 doses only	112 (46.7)	408 (46.4)
5 doses	39 (16.3)	100 (11.4)


Table 7b: Comparison of ARV vaccination by recall and documented evidence for Intra Muscular route of vaccine administration

Arv dose completed	Provides document	Information based on recall
	(n =87)	(n=324)
	n %	n %
1 dose only	10 (11.5)	18 (5.6)
2 doses only	5 (5.8)	24 (7.4)
3 doses only	17 (19.5)	118 (36.4)
4 doses only	17 (19.5)	68 (21.0)
5 doses	38 (43.7)	96 (29.6)

Table 7c: Comparison of ARV vaccination by recall and documented evidence for Intra Dermal route of vaccine administration

Arv dose completed	Provides document	Information based on recall
	(n =153)	(n=556)
	n %	n %
1 dose only	16 (10.5)	28 (5.0)
2 doses only	16 (10.5)	65 (11.7)
3 doses only	25 (16.3)	119 (21.4)
4 doses only	96 (62.7)	344 (61.9)

14. Figure 5: Flow chart illustrating the enrolment of participants in the survey



15. Table 8: Socio-demographic characteristics of study participants by regions, Human rabies death and animal bite burden survey, India, 2022-2023

Socio-demographic characteristics	North	North-East	East	West	South
	Number (% of total)				
Age (in years) (N= 337,808)	(N=73,674)	(N=67,787)	(N=65,806)	(N=68,788)	(N=61,753)
0-14	18,291 (24.8)	16,065 (23.7)	16,670 (25.3)	15,448 (22.5)	12,392 (20.1)
15-60	49,754 (67.5)	46,729 (68.9)	44,527 (67.7)	47,594 (69.2)	42,139 (68.2)
>60	5,629 (7.7)	4,993 (7.4)	4,609 (7.0)	5,746 (8.3)	7,222 (11.7)
Median (IQR)	27 (15-42)	28 (15-43)	28 (14-43)	29 (16-45)	33 (18-50)
Gender (N= 337,808)	(N=73,674)	(N=67,787)	(N=65,806)	(N=68,788)	(N=61,753)
Male	38,212 (51.9)	34,241 (50.5)	33,767 (51.3)	35,262 (51.3)	29,862 (48.3)
Female	35,431 (48.1)	33,507 (49.4)	31,999 (48.6)	33,502 (48.7)	31,860 (51.6)
Others	31 (0.04)	39 (0.1)	40 (0.1)	24 (0.03)	31 (0.1)
Education of the head of the household (N=78,807)	(N= 16,010)	(N= 15,494)	(N= 15,515)	(N= 15,920)	(N= 15,868)
Professional or honors/graduate	1,769 (11.0)	1,335 (8.6)	1,763 (11.4)	2,148 (13.5)	2,193 (13.8)
Senior/higher secondary (11 & 12)	2,529 (15.8)	1,637 (10.6)	2,116 (13.6)	1,997 (12.5)	1,784 (11.2)
High school certificate (9 & 10)	3,570 (22.3)	4,093 (26.4)	2,740 (17.7)	3,874 (24.3)	3,964 (25.0)
Middle school certificate (6-8)	2,528 (15.8)	3,353 (21.6)	2,823 (18.2)	2,958 (18.6)	2,566 (16.2)
Primary school certificate (1-5)	2,610 (16.3)	2,616 (16.9)	2,909 (18.7)	2,873 (18.0)	2,362 (14.9)
Illiterate/ read and write with no formal education	3,004 (18.8)	2,460 (15.9)	3,164 (20.4)	2,070 (13.0)	2,999 (18.9)

Socio-demographic characteristics	North	North-East	East	West	South
	Number (% of total)				
Occupation of the head of the household (N=78,807)	(N= 16,010)	(N= 15,494)	(N= 15,515)	(N= 15,920)	(N= 15,868)
Professional/managers/officials/legislator	313 (2.0)	299 (1.9)	258 (1.7)	356 (2.2)	521 (3.3)
Technicians/associate professionals	438 (2.7)	385 (2.5)	277 (1.8)	944 (5.9)	675 (4.2)
Clerks/clerical support workers	734 (4.6)	509 (3.3)	644 (4.2)	781 (4.9)	700 (4.4)
Service and sales workers	5,127 (32.0)	3,167 (20.4)	3,171 (20.4)	3,162 (19.9)	3,506 (22.1)
Agricultural and fishery workers	3,742 (23.4)	4,803 (31.0)	7,180 (46.3)	7,162 (45.0)	4,693 (29.6)
Craft and related trade workers	3,808 (23.8)	3,970 (25.6)	2,653 (17.1)	2,265 (14.2)	3,373 (21.3)
Homemaker	487 (3.0)	887 (5.7)	480 (3.1)	402 (2.5)	1,098 (6.9)
Unemployed	675 (4.2)	545 (3.5)	477 (3.1)	293 (1.8)	601 (3.8)
Retired	686 (4.3)	929 (6.0)	375 (2.4)	555 (3.5)	701 (4.4)
Type of house (N= 78,807)	(N= 16,010)	(N= 15,494)	(N= 15,515)	(N= 15,920)	(N= 15,868)
Pucca	12,243 (76.5)	3,838 (24.8)	6,063 (39.1)	8,599 (54.0)	11,341 (71.5)
Semi-pucca	2,945 (18.4)	3,930 (25.3)	6,122 (39.4)	4,268 (26.8)	3,430 (21.6)
Kuccha	822 (5.1)	7,726 (49.9)	3,330 (21.5)	3,053 (19.2)	1,097 (6.9)
Toilet facility (N= 78,807)	(N= 16,010)	(N= 15,494)	(N= 15,515)	(N= 15,920)	(N= 15,868)
Own toilet (outside or inside the house)	12,451 (77.8)	11,318 (73.0)	10,489 (67.6)	12,666 (79.6)	14,364 (90.5)
Shared common toilet	2,215 (13.8)	3,745 (24.2)	2,736 (17.6)	1,309 (8.2)	600 (3.8)
Public toilet	17 (0.1)	64 (0.4)	716 (4.6)	183 (1.1)	159 (1.0)
Use open space	1,327 (8.3)	367 (2.4)	1,574 (10.1)	1,762 (11.1)	745 (4.7)
Any member in the household received pre-exposure prophylaxis (N=78,807)	(N= 16,010)	(N= 15,494)	(N= 15,515)	(N= 15,920)	(N= 15,868)
Received	61 (0.4)	47 (0.3)	13 (0.1)	165 (1.0)	167 (1.1)

16. Table 9: Characteristics of Animal Bite among study participants by regions, Human rabies death and animal bite burden survey, India, 2022-2023

Characteristics of animal and dog bite	North	North-East	East	West	South
	Number	Number	Number	Number	Number
Animal bites reported in the last 5 years	1,245	989	683	1,116	1,276
Dog bites reported in the last 5 year	1,148	670	602	1,063	806
Animal bites reported in the last 3 years	914	803	497	831	1,028
Dog bites reported in the last 3 year	835	523	436	788	636
Animal bites reported in the last 1 year	457	431	262	385	517
Dog bites reported in the last 1 year	409	261	227	359	320
Annual incidence per 1000 population	Incidence [95% CI]				
Unweighted annual incidence of animal bite	6.2 [5.6-6.8]	6.36 [5.78-7.0]	4.0 [3.5-4.5]	5.6 [5.0-6.2]	8.2 [7.7-9.1]
Weighted annual incidence of animal bite	7.5 [5.7-9.9]	7.8 [6.3-9.6]	4.4 [3.6-5.2]	5.5 [4.7-6.4]	9.7 [8.2-11.6]
Unweighted annual incidence of dog bite	5.5 [5.0-6.1]	3.8 [3.4-4.3]	3.4 [3.0-3.9]	5.2 [4.7-5.8]	5.2 [4.6-5.8]
Weighted annual incidence of dog bite	6.5 [4.8-8.8]	4.9 [3.7-6.3]	3.8 [3.1-4.7]	5.0 [4.2-5.9]	7.6 [6.0-9.5]

Characteristics of animal and dog bite	North	North-East	East	West	South
	Incidence [95% CI]				
Weighted annual incidence of dog bite by age (in years) of the bite victim					
0-14	8.0 [4.9-13.2]	6.6 [4.5-9.8]	6.0 [4.5-7.9]	6.9 [5.6-8.6]	10.5 [7.6-14.4]
15-60	5.5 [4.3-6.9]	4.2 [3.0-5.7]	2.8 [2.1-3.6]	4.3 [3.5-5.4]	6.6 [5.2-8.4]
>60	10.7 [4.9-23.8]	5.6 [2.7-11.5]	6.0 [3.7-9.7]	5.8 [3.9-8.6]	8.2 [5.4-12.4]
Weighted annual incidence of dog bite by gender of the bite victim					
Male	9.1 [6.8-12.0]	6.0 [4.9-7.4]	4.9 [4.0-6.1]	6.9 [5.8-8.3]	8.8 [6.4-12.2]
Female	3.6 [2.2-6.1]	3.6 [2.2-6.0]	2.7 [1.9-3.7]	3.0 [2.2-3.9]	6.4 [4.8-8.4]
Weighted annual incidence of dog bite by type of residence					
Rural	6.8 [4.9-9.6]	4.8 [3.6-6.3]	3.4 [2.7-4.3]	5.0 [4.1-6.0]	8.4 [6.5-11.0]
Urban	4.1 [2.9-5.8]	6.1 [3.0-12.7]	6.0 [4.2-8.7]	5.0 [3.5-7.2]	5.7 [3.7-8.8]

Characteristics of animal and dog bite	North	North-East	East	West	South
	Number (% of Total)				
Biting animal (n= 2,052)	(n=457)	(n=431)	(n=262)	(n=385)	(n=517)
Dog	409 (89.5)	261 (60.6)	227 (86.6)	359 (93.2)	320 (61.9)
Cat	17 (3.7)	160 (37.1)	28 (10.7)	15 (3.9)	165 (31.9)
Monkey	20 (4.4)	3 (0.7)	5 (1.9)	8 (2.1)	12 (2.3)
Rat	6 (1.3)	4 (0.9)	1 (0.4)	-	14 (2.7)
Others (donkey, cow, domestic pig, rabbit, jackal, leopard)	5 (1.1)	3 (0.6)	1 (0.4)	3(0.8)	6 (1.2)
Category of dog (n= 1,576)	(n=409)	(n=261)	(n=227)	(n=359)	(n=320)
Pet	149 (36.4)	151 (57.8)	43 (18.9)	83 (23.1)	188 (58.7)
Stray	260 (63.6)	110 (42.2)	184 (81.1)	276 (76.9)	132 (41.3)
Nature of attack (n= 1,576)	(n=409)	(n=261)	(n=227)	(n=359)	(n=320)
Unprovoked attack	339 (82.9)	206 (78.9)	175 (77.1)	305 (85.0)	256 (80.0)
Provoked attack	70 (17.1)	55 (21.1)	52 (22.9)	54 (15.0)	64 (20.0)
Availability of the dog for 10 days (n=1,576)	(n=409)	(n=261)	(n=227)	(n=359)	(n=320)
Available	335 (81.9)	199 (76.2)	123 (54.2)	253 (70.5)	234 (73.1)
Status of the dog after 10 days (n=1,144)	(n=335)	(n=199)	(n=123)	(n=253)	(n=234)
Alive	306 (91.3)	193 (97.0)	113 (91.9)	236 (93.3)	218 (93.2)
Died	11 (3.3)	2 (1.0)	3 (2.4)	8 (3.2)	7 (3.0)
Killed within 10 days	4 (1.2)	3 (1.5)	7 (5.7)	3 (1.2)	5 (2.1)
Don't know	14 (4.2)	1 (0.5)	-	6 (2.4)	4 (1.7)

Characteristics of animal and dog bite	North	North-East	East	West	South
	Number (% of Total)				
Number of wounds (n= 1,576)	(n=409)	(n=261)	(n=227)	(n=359)	(n=320)
One	381 (93.2)	249 (95.4)	207 (91.2)	315 (87.7)	305 (95.3)
Two	18 (4.4)	10 (3.8)	18 (7.9)	31 (8.6)	11 (3.4)
More than two	10 (2.4)	2 (0.8)	2 (0.9)	13 (3.6)	4 (1.3)
Site of bite (n= 1,734)	(n=451)	(n=275)	(n=250)	(n=418)	(n=340)
Leg/feet	326 (72.3)	171 (62.2)	174 (69.6)	288 (68.9)	223 (65.6)
Arm/forearm/hands	109 (24.2)	77 (28.0)	57 (22.8)	88 (21.1)	97 (28.5)
Trunk	10 (2.2)	11 (4.0)	14 (5.6)	28 (6.7)	12 (3.5)
Head/face	6 (1.3)	16 (5.8)	5 (2.0)	14 (3.3)	8 (2.4)
Type of wound (n= 1,734)	(n=451)	(n=275)	(n=250)	(n=418)	(n=340)
Abrasion	265 (58.8)	177 (64.4)	167 (66.8)	288 (68.9)	233 (68.5)
Laceration	138 (30.6)	80 (29.0)	45 (18.0)	62 (14.8)	69 (20.3)
Puncture wound	43 (9.5)	17 (6.2)	38 (15.2)	62 (14.8)	36 (10.6)
Avulsion	5 (1.1)	1 (0.4)	0 (0.0)	6 (1.5)	2 (0.6)
Wound management (n=1,576)	(n=409)	(n=261)	(n=227)	(n=359)	(n=320)
Wound wash done	286 (69.9)	209 (80.1)	144 (63.4)	188 (52.4)	218 (68.1)
Locally available applicant	128 (31.3)	6 (2.3)	23 (10.1)	61 (17.0)	58 (18.1)
Applied antiseptic	43 (10.5)	104 (39.8)	44 (19.4)	64 (17.8)	29 (9.1)

Characteristics of animal and dog bite	North	North-East	East	West	South
	Number (% of Total)				
Use to wash the wound(s) (n=1,045)	(n=286)	(n=209)	(n=144)	(n=188)	(n=218)
Wash with water and soap	254 (88.8)	164 (78.5)	130 (90.3)	151 (80.3)	150 (68.8)
Wash with water alone	32 (11.2)	45 (21.5)	14 (9.7)	37 (19.7)	68 (31.2)
Locally available applicants (n=276)	(n=128)	(n=6)	(n=23)	(n=61)	(n=58)
Chilly powder	86 (67.2)	-	1	19 (31.1)	3 (5.2)
Turmeric powder	29 (22.7)	1	9	29 (47.5)	39 (67.2)
Others (snuff, powdered tablet, oil, salt)	7 (5.4)	2	12	13 (21.4)	8 (13.8)
Plant sap/Coconut water	6 (4.7)	2	1	-	1 (1.7)
Others (Mud, Coffee Powder, Cow Dung, Coin, Snuff, Powdered tablet, Oil, Salt)		1			7 (12.1)
Medical care at a health facility (n=1,576)	(n=409)	(n=261)	(n=227)	(n=359)	(n=320)
Yes	334 (81.7)	191 (73.2)	200 (88.1)	285 (79.4)	307 (95.9)
Time gap between the bite and medical care sought (n=1,317)	(n=334)	(n=191)	(n=200)	(n=285)	(n=307)
Within 1 day	244 (73.1)	159 (83.2)	158 (79.0)	255 (89.4)	251 (81.8)
1 day - 3 days	85 (24.4)	29 (15.2)	37 (18.5)	29 (10.2)	56 (18.2)
3 days - 7 days	4 (1.2)	3 (1.6)	1 (0.5)	-	-
More than 7 days	1 (0.3)	-	4 (2.0)	1 (0.3)	-

Characteristics of animal and dog bite	North	North-East	East	West	South
	Number (% of Total)				
Post-exposure prophylaxis coverage of dog bite victims					
Received anti rabies vaccine (ARV) (n=1,576)	314 (76.8)	174 (66.7)	194 (85.5)	272 (75.8)	299 (93.4)
Type of health facility (n=1,253)	(n=314)	(n=174)	(n=194)	(n=272)	(n=299)
Public health facility (PHC/CHC, DH/SDH/TH, MCH)	261 (83.1)	131 (75.3)	185 (95.4)	255 (93.7)	289 (96.7)
Private health facility	53 (16.9)	43 (24.7)	9 (4.6)	17 (6.3)	10 (3.3)
Route of administration (n=1,253)	(n=314)	(n=174)	(n=194)	(n=272)	(n=299)
Intra dermal (ID)	244 (77.7)	57 (32.8)	102 (52.6)	239 (87.9)	194 (64.9)
Intra muscular (IM)	70 (22.3)	117 (67.2)	92 (47.4)	33 (12.1)	105 (35.1)
Site of vaccination (n=1,253)	(n=314)	(n=174)	(n=194)	(n=272)	(n=299)
Deltoid	305 (97.1)	166 (95.4)	190 (97.9)	261 (96.0)	295 (98.7)
Antero lateral thigh	1 (0.3)	6 (3.4)	1 (0.5)	2 (0.7)	1 (0.3)
Gluteal	8 (2.6)	2 (1.2)	3 (1.6)	9 (3.3)	3 (1.0)

Characteristics of animal and dog bite	North	North-East	East	West	South
	Number (% of Total)				
ARV dose completed (ID route of administration) (n=836)	(n=244)	(n=57)	(n=102)	(n=239)	(n=194)
1 dose only	19 (7.8)	2 (3.5)	5 (4.9)	17 (7.1)	12 (6.2)
2 doses only	27 (11.1)	5 (8.8)	7 (6.9)	38 (15.9)	19 (9.8)
3 doses only	69 (28.3)	9 (15.8)	9 (8.8)	51 (21.3)	43 (22.2)
4 doses	129 (52.8)	41 (71.9)	81 (79.4)	133 (55.7)	120 (61.8)
ARV dose completed (IM route of administration) (n=417)	(n=70)	(n=117)	(n=92)	(n=33)	(n=105)
1 dose only	9 (12.9)	7 (6.0)	2 (2.2)	-	11 (10.5)
2 doses only	3 (4.3)	2 (1.7)	8 (8.7)	5 (15.2)	12 (11.4)
3 doses only	30 (42.8)	31 (26.5)	39 (42.4)	8 (24.2)	29 (27.6)
4 doses only	9 (12.9)	10 (8.5)	12 (13.0)	4 (12.1)	52 (49.5)
5 doses	19 (27.1)	67 (57.3)	31 (33.7)	16 (48.5)	1 (1.0)

Characteristics of animal and dog bite	North	North-East	East	West	South
	Number (% of Total)				
Reason for not vaccinated (n=323)	(n=95)	(n=87)	(n=33)	(n=87)	(n=21)
Not aware of post exposure prophylaxis/ Don't know where to get the vaccine	2 (2.1)	1 (1.2)	-	1 (1.2)	2
Generally, don't believe in vaccine/Vaccine will not be helpful in preventing the disease	54 (56.8)	53 (60.9)	25 (75.8)	41 (47.1)	10
Consulted a traditional healer/Siddha/Ayurveda practitioner	-	9 (10.4)	-	-	1
Minor scratch	2 (2.1)	3 (3.4)	-	2 (2.3)	-
Can't miss work	1 (1.1)	6 (6.9)	-	6 (6.9)	2
Only TT taken	2 (2.1)	2 (2.3)	-	2 (2.3)	1
Couldn't afford the vaccine	17 (17.9)	3 (3.4)	-	7 (8.0)	1
Bitten by a vaccinated dog	8 (8.4)	2 (2.3)	-	2 (2.3)	1
Hospital too far/No transportation	4 (4.2)	2 (2.3)	2 (6.1)	9 (10.3)	2
Fear of side effects, pain at the injection site and fever/ Heard from social media/neighbour/religious leader/mass media that vaccine is harmful	2 (2.1)	2 (2.3)	3 (9.1)	16 (18.4)	1
Vaccine not available	3 (3.2)	4 (4.6)	1 (3.0)	1 (1.2)	-

Characteristics of animal and dog bite	North	North-East	East	West	South
	Number (% of Total)				
Received rabies immunoglobulin (RIG) (N=1,576)	7 (1.7)	1 (0.4)	13 (5.7)	6 (1.7)	54 (16.9)
Site of RIG (N=81)	(n=7)	(n=1)	(n=13)	(n=6)	(n=54)
Into the wound	5	1	10	5	27 (50.0)
Into the wound and deltoid	1	-	3	1	23 (42.6)
Into the wound & gluteal	1	-	-	-	1 (1.9)
Into the wound & antero lateral thigh	-	-	-	-	3 (5.6)
Type of health facility RIG was received (N=81)					
Public health facility (PHC/CHC, DH/SDH/TH, MCH)	6	-	11	5	50 (92.6)
Private Health Facility	1	1	2	1	4 (7.4)
Type of wound received RIG					
Avulsion (n=11)	-	1	-	1	1
Puncture (n=179)	4 (2.2)	-	7 (3.9)	4 (2.2)	15 (8.5)
Laceration (n=356)	3 (0.8)	-	4 (1.2)	1 (0.3)	20 (5.6)
Abrasion (n=1,030)	-	-	2 (0.2)	-	18 (1.7)
Documentation for vaccination (n=1,120)	(n=314)	(n=174)	(n=194)	(n=272)	(n=166)
Provides information based on recall	237 (75.5)	108 (62.1)	162 (83.5)	234 (86.0)	139 (83.7)
Provides documentation	77 (24.5)	66 (37.9)	32 (16.5)	38 (14.0)	27 (16.3)

17. Table 10: Household pet ownership and rabies vaccination by regions, Human rabies death and animal bite burden survey, India, 2022-2023

Characteristics of the pet/domestic animal	North	North-East	East	West	South
	Number (% of total)				
	(n=16,010)	(n=15,494)	(n= 15,515)	(n=15,920)	(n=15,868)
Owns pet/domestic animal	4,111 (25.7)	5,923 (38.2)	3,252 (21.0)	4,484 (28.2)	2,224 (14.0)
Type of pet (n=19,994)	(n=4,111)	(n=5,923)	(n=3,252)	(n=4,484)	(n=2,224)
Dog	846 (20.6)	1,301 (22)	196 (6)	351 (7.8)	1,020 (45.9)
Cat	71 (1.7)	1,677 (28.3)	135 (4.2)	307 (6.8)	627 (28.2)
Cow	2,062 (50.2)	2,585 (43.6)	2,680 (82.4)	2,395 (53.4)	509 (22.9)
Goat/sheep	934 (22.7)	1,707 (28.8)	618 (19.0)	1,257 (28)	621 (27.9)
Pig	5 (0.1)	1,611 (27.2)	14 (0.4)	2 (0.04)	5 (0.2)
Horse	17 (0.4)	3 (0.1)	1 (0.03)	8 (0.2)	2 (0.1)
Donkey	6 (0.1)	1 (0.02)	-	3 (0.1)	-
Rabbit	37 (0.9)	58 (1.0)	7 (0.2)	8 (0.2)	21 (0.9)
Monkey	-	6 (0.1)	2 (0.1)	-	-
Others (buffalo, ox, camel, mongoose)	1,125 (27.4)	7 (0.1)	217 (6.7)	1,571 (35)	46 (2.1)
No of pet dogs in each household (n=3,714)	(n=846)	(n=1,301)	(n=196)	(n=351)	(n=1,020)
One dog	774 (91.5)	1,042 (80.0)	182 (92.9)	317 (90.3)	871 (85.4)
Two dogs	59 (7.0)	188 (14.5)	12 (6.1)	22 (6.3)	120 (11.8)
Three or more dogs	13 (1.5)	71 (5.5)	2 (1.0)	12 (3.4)	29 (2.8)

Characteristics of the pet/domestic animal	North	North-East	East	West	South
	Number (% of total)				
Vaccination status of the dog (n=4,369)	(n=931)	(n=1,631)	(n=212)	(n=397)	(n=1,198)
Vaccinated	612 (65.7)	551 (33.8)	93 (43.9)	186 (46.9)	781 (65.2)
Not vaccinated	289 (31.1)	1057 (64.8)	114 (53.8)	195 (49.1)	397 (33.1)
Don't know about vaccination status	30 (3.2)	23 (1.4)	5 (2.3)	16 (4.0)	20 (1.7)
Own dog(s) died in the past year (N=78,807)	(n=16,010)	(n=15,494)	(n= 15,515)	(n=15,920)	(n=15,868)
Yes	261 (1.6)	275 (1.8)	74 (0.5)	82 (0.5)	458 (2.9)
Provide care for other dogs (N=78,807)	(n=16,010)	(n=15,494)	(n= 15,515)	(n=15,920)	(n=15,868)
Stray dog	6,748 (42.2)	714 (4.6)	1,522 (9.8)	3,418 (21.5)	813 (5.1)
Community dog	420 (2.6)	861 (5.6)	503 (3.2)	5,153 (32.4)	280 (1.8)
Does not provide care	8,842 (55.2)	13,919 (89.8)	13,490 (87.0)	7,349 (46.1)	14,775 (93.1)
Type of care provide for the stray/community dogs (n=20,432)	(n=7,168)	(n=1,575)	(n= 2,025)	(n=8,571)	(n=1,093)
Food	6,671 (93.1)	1,484 (94.2)	2,014 (99.5)	8,532 (99.5)	1,058 (96.8)
Water	5,944 (82.9)	1,150 (73)	1,488 (73.5)	5,093 (59.4)	658 (60.2)
Shelter	232 (3.2)	208 (13.2)	142 (7.0)	63 (0.7)	308 (28.2)
Veterinary care	7 (0.1)	1 (0.1)	1 (0.1)	2 (0.02)	6 (0.5)

18. Table 11: Comparison of the age, gender and type of residence of the participants in the study with that of the Census 2011 India population

India	Rabies study population (N=337,808)	2011 Census Population (N=2,057,220,656)
Age (in Years)	n (%)	n (%)
0-5	27,256 (8.1)	137,406,673 (13.4)
6-18	78,147 (23.1)	313,088,772 (30.4)
19-60	204,206 (60.4)	516,896,063 (50.3)
>60	28,199 (8.4)	58,480,348 (5.7)
Gender		
Male	171,344 (50.7)	1,064,313,544 (51.7)
Female	166,299 (49.2)	992,907,112 (48.3)
Type of residence		
Rural	239,095 (70.8)	1,484,981,278 (72.2)
Urban	98,713 (29.2)	572,239,378 (27.8)

19. STROBE Statement—Checklist of items that should be included in reports of cross-sectional studies

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	3
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4
Objectives	3	State specific objectives, including any prespecified hypotheses	4
Methods			
Study design	4	Present key elements of study design early in the paper	6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6,7
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	7
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	7
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	NA
Bias	9	Describe any efforts to address potential sources of bias	6,7
Study size	10	Explain how the study size was arrived at	6,7
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	8
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	7,8
		(b) Describe any methods used to examine subgroups and interactions	7,8
		(c) Explain how missing data were addressed	NA
		(d) If applicable, describe analytical methods taking account of sampling strategy	7,8
		(e) Describe any sensitivity analyses	8
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Apx Figure 2
		(b) Give reasons for non-participation at each stage	Apx Figure 2
		(c) Consider use of a flow diagram	Apx Figure 2

Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Table 1 Page No 8,9
		(b) Indicate number of participants with missing data for each variable of interest	Na
Outcome data	15*	Report numbers of outcome events or summary measures	Table 2,3, Page No 9,10
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 2, Page No 9,10
		(b) Report category boundaries when continuous variables were categorized	Table 1, Page No 9,10
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	NA
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Apx Table 4, Page no 9,10
Discussion			
Key results	18	Summarise key results with reference to study objectives	11,
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	14
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	11-14
Generalisability	21	Discuss the generalisability (external validity) of the study results	11
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	8

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.



Estimates of the burden of human rabies deaths and animal bites in India, 2022–23: a community-based cross-sectional survey and probability decision-tree modelling study

Jeromie Wesley Vivian Thangaraj, Navaneeth S Krishna, Shanmugasundaram Devika, Suganya Egambaram, Sudha Rani Dhanapal, Siraj Ahmed Khan, Ashok Kumar Srivastava*, Ayush Mishra*, Basavaraj Shrinivasa*, Devendra Gour*, Major Madhukar*, Nirmal Verma*, Parul Sharma*, Ravinder Kumar Soni*, Sabarinathan Ramasamy*, Sreelakshmi Mohandas*, Subrata Baidya*, Tanveer Rehman*, Vijay V Yeldandi*, Akashdeep Singh†, Aswathy Sreedevit, Babasaheb V Tandale†, Debjani Ram Purakayastha†, Mahendra M Reddy†, Manju Toppot, Nitinkumar Valjibhai Solankit, Pramit Ghosh†, Prashant Jaiswal†, Shaili Vyast, Shampa Das†, Subrata Kumar Palot, Venela Prasant†, Amanda G A Rozario, Chokkalingam Durairajan, Anitha Delli, Aruna Sasi, Chandhini Pandiyan, Doddabale Hanumanthaiiah Ashwathnarayana, Sam Joy, Srikrishna Isloor, Mysore Kalappa Sudarshan, Manju Rahi, Manoj V Murhekar, and Human Rabies Study Collaborators‡

Summary

Background Reliable and recent data of human rabies deaths and animal bites are not available in India, where a third of global cases occur. Since there is a global target of eliminating dog-mediated human rabies by 2030, understanding whether the country is on track is essential. We aimed to estimate the animal-bite burden and the number of human rabies deaths in India.

Methods We conducted a community-based nationwide cross-sectional survey with a multistage cluster-sampling design from March 2, 2022 to Aug 26, 2023, covering 60 districts in 15 Indian states. The head of the household or an adult family member was interviewed to collect information about animal-bite history in family members, receipt of anti-rabies vaccination (ARV), and death following animal bite in the family. Annual animal-bite incidence along with 95% CIs were estimated after applying the sampling weights and adjusting for clustering. We estimated annual human rabies deaths using a decision-tree probability model with parameters from the community survey and laboratory data on rabies positivity among suspected rabid dogs.

Findings Of the 337 808 individuals residing in the 78 807 households surveyed, 2052 gave a history of animal bite, mostly (1576 [76·8%]) due to dogs in the past 1 year. The weighted and adjusted annual incidence of animal bite was 6·6 (95% CI 5·7–7·6) per 1000 population, translating into 9·1 million bites nationally. Annual dog-bite incidence was 5·6 (4·8–6·6) per 1000. Among people who had been bitten by a dog, 323 (20·5%) did not receive ARV, and 1043 (66·2%) received at least three doses. Nearly half (615 [49·1%]) of the 1253 individuals who received one dose did not complete their full course of vaccination. We estimated 5726 (95% uncertainty interval 3967–7350) human rabies deaths occurring annually in India.

Interpretation Although there was a substantial decline in human rabies deaths over the past two decades, to eliminate dog-mediated human rabies by 2030, India needs to fast-track its actions by adopting a focused one-health approach. Integrating human and animal surveillance, ensuring timely administration of full course of post-exposure prophylaxis, and accelerating dog vaccination across the country are crucial steps towards this goal.

Funding Indian Council of Medical Research.

Copyright © 2024 Elsevier Ltd. All rights reserved, including those for text and data mining, AI training, and similar technologies.

Introduction

Rabies, a fatal but vaccine-preventable viral zoonotic disease, causes approximately 59 000 human deaths globally each year, mostly because of dog bites.¹ United Against Rabies, an alliance of partners from international agencies, has made a global call to end human deaths from dog-mediated rabies by 2030 (Zero by 30).² To achieve this goal, India has laid a roadmap through its National Action Plan for Dog-Mediated Rabies Elimination from India by 2030 (NAPRE).³ NAPRE

envisages a one-health approach, involving human and veterinary health sectors, the Agriculture Ministry, municipal corporations, and Panchayati Raj Institutions to achieve 75% dog-vaccination coverage by 2025, and 75% reduction in human rabies deaths by 2030. The key component of the action plan includes mass dog vaccination, ensuring availability of rabies biologicals free of cost at all levels of the health system and strengthening linkages between rabies surveillance in humans and animals.

Lancet Infect Dis 2025;

25: 126–34

Published Online

September 30, 2024

[https://doi.org/10.1016/S1473-3099\(24\)00490-0](https://doi.org/10.1016/S1473-3099(24)00490-0)

See Comment page 13

This online publication has been corrected. The corrected version first appeared at thelancet.com/infection on December 27, 2024

*Contributed equally

†Contributed equally

‡Members listed at the end of the Article

Indian Council of Medical Research (ICMR)-National Institute of Epidemiology, Chennai, India

(J WV Thangaraj MPH,

N S Krishna MBBS, S Devika PhD,

S Egambaram MSc,

S R Dhanapal MPH,

S Ramasamy BE,

A G A Rozario MSc,

C Durairajan MCA, A Delli MSc,

A Sasi MPH, C Pandiyan MPH,

M V Murhekar MD); ICMR-

Regional Medical Research

Centre, Dibrugarh, India

(P Ghosh MD, S A Khan PhD);

Himalayan Institute of Medical

Sciences, Swami Rama

Himalayan University,

Dehradun, India

(A K Srivastava MD, S Vyas MD);

ICMR-Regional Medical

Research Centre, Gorakhpur,

India (A Mishra MBBS,

M M Reddy MD); ICMR-National

Institute of Virology, Pune,

India (B Shrinivasa PhD,

B V Tandale MD); Gandhi

Medical College, Bhopal, India

(D Gour MD, M Toppo MD);

ICMR-Rajendra Memorial

Research Institute of Medical

Sciences, Patna, India

Research in context

Evidence before this study

We searched PubMed and Embase on June 11, 2024, for articles reporting estimates on human rabies deaths and animal-bite incidence in India, using search terms such as “burden”, “incidence”, “mortality”, and “death” in combination with “rabies”, “animal bite”, “dog bite”, and “India”, published in English since Jan 1, 2004. We identified four studies related to human rabies deaths estimation; two modelling studies done in 2005 and 2015, one verbal autopsy study for deaths that occurred in 2005, and a community-based survey done in 2003. All the studies estimated around 20 000 human rabies deaths per year except for the verbal autopsy study, which reported 12 700 deaths. Additionally, we identified 40 articles reporting animal or dog-bite incidence. Of these, 31 were hospital-based studies, and nine were community-based studies. A systematic review done in 2021 reported crude annual dog-bite incidence to be between 0.26% and 2.50%. However, all the studies were done in small geographical areas, using convenient sampling methods within the catchment area of tertiary-care hospitals and did not have generalisability. Human rabies surveillance data in India are unreliable. In this context, we did a nationwide community survey to estimate animal-bite burden, human rabies deaths, and post-exposure prophylaxis (PEP) coverage.

Added value of this study

Current estimates on human rabies death burden, animal-bite burden, and PEP coverage are crucial for India to prioritise strategies to achieve the global goal of eliminating dog-mediated human rabies by 2030 (the Zero by 30 goal).

In India, 17.4 million animal bites occur annually, resulting in 20 565 human rabies deaths.^{4,5} A multicentric community survey on animal bites done in seven states in 2017 demonstrated an anti-rabies vaccination (ARV) coverage of 79.6%.⁶ An evaluation of the human rabies surveillance system indicated a substantial under-reporting of the number of bites and human rabies death burden.⁷ The last nationwide survey to estimate human rabies deaths and animal-bite burden was done in 2003, and this type of up-to-date country-specific data are necessary for reliable global estimates.⁸ In this context, we did a community-based cross-sectional survey across India to estimate animal-bite incidence. Using relevant parameters from the survey, we estimated the number of human rabies deaths using a probability decision-tree model.

Methods

Study design and participants

India has 788 districts across 28 states and eight union territories. We did a community-based nationwide cross-sectional survey in 60 districts across 15 Indian states between March 2, 2022, and Aug 26, 2023 using a

A population-based nationwide survey to obtain information on bite burden and human rabies deaths has not been done in the past two decades. Our survey spanned different regions of India, covering a wide geographical area to ensure representative estimates at national and regional levels. Our study findings indicated that around 9.1 million animal bites occur every year in India, with the majority caused by dogs. Although 80% of the dog-bite victims received one dose of anti-rabies vaccination, only two-thirds received three doses, and 40% completed the prescribed vaccination schedule. Only a tenth of the eligible bite victims received passive immunisation. Our survey also found that around half of the pet dogs in the households were not vaccinated against rabies. We used the information from the survey and animal laboratories and estimated 5726 human rabies deaths per year using the probability decision-tree model, showing nearly a 75% decrease in the number of deaths in the past 20 years.

Implications of all the available evidence

Newer estimates will have substantial implications on the global burden of rabies, given that India contributes around 35% of the global human rabies death burden. Findings from this study will provide clarity on the current status of this burden and enable the programme managers to prioritise their strategies and tailor their efforts to eliminate dog-mediated rabies by 2030. Specifically, India needs to adopt a one-health approach by integrating human and animal surveillance, ensuring timely administration of a full course of PEP for all bite victims, and prioritising scale-up of mass dog vaccination across the country.

multistage cluster-sampling design. For sampling purposes, we divided India into five geographical regions (north, south, east, west, and northeast). We selected three states from each geographical region using simple random sampling. In each state, we selected four districts using the probability proportional-to-size (PPS) method. Within each district, we selected ten clusters (villages in rural areas and wards in urban areas) using the PPS method. From each cluster, we randomly selected one census enumeration block (geographical area within a village or ward delineated as part of decadal census with around 130–150 households; figure 1; appendix p 1).

Assuming an animal-bite incidence of 1.7%,⁵ relative margin of error of 10%, design effect of 2%, confidence level of 95%, and non-response of 15%, we required a sample size of 52 267 people per geographical region. With a minimum household size of 3.5 (census 2011), we needed to cover approximately 15 000 households (around 5000 households per state) in each region. Therefore, a minimum of 140 households were to be visited in each cluster to reach the sample size.

In each selected cluster, the study team visited 140 households sequentially after randomly selecting the

(M Madhukar DTCO, D R Purakayastha PhD); Pt Jawahar Lal Nehru Memorial Medical College, Raipur, India (N Verma MD, P Jaiswal MD); GMERS Medical College and Hospital, Patan, India (P Sharma MD, NV Solanki MD); Dayanand Medical College and Hospital, Ludhiana, India (R K Soni PhD, A Singh MD); Amrita Institute of Medical Sciences, Amrita Vishwa Vidyapeetham, Kochi, India (S Mohandas MD, A Sreedevi MD); Agartala Government Medical College and Model Rural Health Research Unit, Tripura, India (S Baidya MD, S Das MD); ICMR-Regional Medical Research Centre, Bhubaneswar, India (T Rehman MD, S K Palo MD); Society for Health Allied Research and Education, India, Hyderabad, India (V V Yeldandi MD, V Prasanth MBA); Kempegowda Institute of Medical Sciences, Bengaluru, India (D H Ashwathnarayana MD); ICMR, New Delhi, India (S Joy PhD); Karnataka Veterinary, Animal, and Fisheries Sciences University, Veterinary College, Bengaluru, India (S Isloor PhD); Association for the Prevention and Control of Rabies in India, Bengaluru, India (M K Sudarshan MD); ICMR-Vector Control Research Centre, Pondicherry, India (M Rahi MD)

Correspondence to: Dr Manoj V Murhekar, Indian Council of Medical Research-National Institute of Epidemiology, Chennai 600077, India (mmurhekar@nieicmr.org.in)

See Online for appendix




Figure 1: Sampling strategy, study districts
(A) Multistage cluster-sampling design employed for the survey. (B) Map of the 60 selected study districts from 15 states across five regions in India. The map was created in QGIS version 3.34. The shape file was adapted by permission of the Survey of India.

first household. If the selected census enumeration block did not have 140 households, households in the adjacent census enumeration block were included. Trained investigators, using Open Data Kit software on a tablet, collected demographic details and histories of animal bites and deaths caused by animal bite over the past 5 years for all the usual family members residing in the household. Additional information including the type of biting animal, status of the animal after the bite, circumstances leading to the animal bite, and receipt of post-exposure prophylaxis (PEP) was collected for individuals reporting history of animal bites. Dog bites

Study participants	
Age, in years (n=337 808)	
0-14	78 866 (23.4%)
15-60	230 743 (68.3%)
>60	28 199 (8.3%)
Median (IQR)	28 (15-45)
Gender (n=337 808)	
Male	171 344 (50.7%)
Female	166 299 (49.2%)
Other	165 (0.1%)
Education of the head of the household (n=78 807)	
Professional, or honours or graduate degree	9208 (11.7%)
Senior or higher secondary (grade 11 and 12)	10 063 (12.8%)
High-school certificate (grade 9 and 10)	18 241 (23.1%)
Middle-school certificate (grade 6-8)	14 228 (18.0%)
Primary-school certificate (grade 1-5)	13 370 (17.0%)
Illiterate, or can read and write with no formal education	13 697 (17.4%)
Occupation of the head of the household (n=78 807)	
Professional, managers, officials, and legislators	1747 (2.2%)
Technicians and associate professionals	2719 (3.4%)
Clerks or clerical support workers	3368 (4.3%)
Service and sales workers	18 133 (23.0%)
Agricultural and fishery workers	27 580 (35.0%)
Craft and related trade workers	16 069 (20.4%)
Homemaker	3354 (4.3%)
Unemployed	2591 (3.3%)
Retired	3246 (4.1%)
Type of house (n=78 807)	
Pucca	42 084 (53.4%)
Semi-pucca	20 695 (26.3%)
Kuccha	16 028 (20.3%)
Toilet facility (n=78 807)	
Own toilet (outside or inside the house)	61 288 (77.8%)
Shared common toilet	10 605 (13.5%)
Public toilet	1139 (1.4%)
Use open space	5775 (7.3%)

Table 1: Sociodemographic characteristics of study participants

identified during the survey were classified as bite by a normal or suspected rabid dog using the criteria proposed by Tepsumethanon and colleagues.⁹

Institutional Human Ethics Committee of the Indian Council of Medical Research–National Institute of Epidemiology and participating institutions approved the study protocol. We obtained written informed consent from participants older than 18 years, parental consent from those younger than 17 years, and assent from participants aged 7–17 years.

Statistical analysis

We used proportions, mean (SD), and median (IQR) to describe household and participant characteristics. We computed overall weighted annual incidence of animal

and dog bites along with 95% CIs using design weights and adjusting for clustering. We calculated the product of probability of selection of sampling units at each stage and took its inverse to calculate the design weight. Design weight was adjusted for the non-response to calculate the final sampling weight (appendix p 2). We also calculated weighted-bite incidence by age groups, gender, and type of residence along with 95% CIs. In addition, we calculated the PEP coverage for ARV by doses, receipt of rabies immunoglobulin, and wound care. We did a cluster-adjusted univariable and multivariable logistic regression analysis to identify factors associated with non-receipt of ARV. Variables with $p < 0.20$ in the univariable analysis were included in the final model.

We used the WHO recommended decision-tree dog-bite probability model to estimate human rabies deaths by using a series of probability steps developed by Cleaveland and colleagues³⁰ and used for estimation of national and global burden.^{11,31} The input parameters from the community survey included annual incidence of suspected rabid dog bites, ARV coverage, and distribution of dog-bite wounds on the body. The data on rabies positivity among suspected rabid dogs was obtained from six veterinary laboratories catering to 12 Indian states. Estimated human rabies deaths were obtained by multiplying the incidence of suspected rabid dog bites from community survey by the probability of death after being bitten by a suspected rabid dog (appendix pp 3–4). We estimated 95% uncertainty interval (UIs) for the estimated human rabies deaths by using a Monte Carlo simulation for 10000 iterations. Furthermore, we did a sensitivity analysis using the lower and upper CIs of animal rabies positivity data and average incidence of suspected rabies dog bite over a 3-year period. The data were analysed using the survey data analysis module in STATA SE version 17.0, and R version 4.2.2.

Role of the funding source

The funder of the study had no role in the study design, data collection, data analysis, data interpretation, or writing of the report.

Results

We surveyed 78 807 (91.2%) of the 86 444 households (the reasons certain households were not included in the survey are presented in the appendix p 5). History of dog bite was collected for 337 808 individuals from these households (appendix p 5). The median age of the survey participants was 28 years (IQR 15–45) years, and 171 344 (50.7%) were men and 166 299 (49.2%) were women. 13 697 (17.4%) heads of household (ie, the person who was usually responsible for the upkeep, maintenance, and made the financial decisions of the household) were illiterate, 42 084 (53.4%) resided in pucca houses, and 73 032 (92.7%) reported that they used either their own, shared, or a public toilet facility (table 1).

2052 individuals reported a history of animal bites in the past year, with 1576 (76.8%) attributed to dogs

	Animal
Animal bites reported in the past 1 year	2052
Dog bites reported in the past 1 year	1576
Biting animal (n=2052)*	
Dog	1576 (76.8%)
Cat	385 (18.8%)
Monkey	48 (2.3%)
Rat	25 (1.3%)
Other (donkey, cow, domestic pig, rabbit, jackal, or leopard)	18 (0.8%)
Category of dog (n=1576)	
Pet	614 (39.0%)
Stray	962 (61.0%)
Nature of attack (n=1576)	
Unprovoked attack	1281 (81.3%)
Provoked attack	295 (18.7%)
Availability of the dog for 10 days (n=1576)	
Available	1144 (72.6%)
Status of the dog after 10 days (n=1144)	
Alive	1066 (93.2%)
Died	31 (2.7%)
Killed within 10 days	22 (1.9%)
Not known	25 (2.2%)
Number of wounds (n=1576)	
One	1457 (92.4%)
Two	88 (5.6%)
More than two	31 (2.0%)
Site of bite (n=1734)	
Lower limbs (leg or feet)	1182 (68.2%)
Upper limbs (arm, forearm, or hands)	428 (24.7%)
Torso	75 (4.3%)
Head or face	49 (2.8%)
Type of wound (n=1734)	
Abrasion	1130 (65.2%)
Laceration	394 (22.7%)
Puncture wound	196 (11.3%)
Avulsion	14 (0.8%)
*Number (% of total).	

Table 2: Characteristics of the biting animal and profile of the bites

(table 2). The annual unweighted animal-bite incidence was 6.1 (95% CI 5.8–6.3) per 1000 population and the annual incidence of dog bite was 4.7 (4.4–4.9) per 1000 population. The weighted incidence after adjusting for clustering was 6.6 (5.7–7.6) for animal bite and 5.6 (4.8–6.6) for dog bite per 1000 per year. Annual dog-bite incidence was 7.5 per 1000 among children aged 0–14 years, 7.8 per 1000 among people older than 60 years, and 7.6 per 1000 among male individuals and 3.5 per 1000 among females. The dog-bite incidence was 5.8 per 1000 in rural areas and 5.0 per 1000 in urban areas (figure 2; appendix p 6). The annual dog-bite incidence ranged between 3.8 to 7.6 per 1000 in different regions (appendix pp 8, 21).




Figure 2: Annual incidence of animal and dog bites overall and by demographic characteristics and post-exposure prophylaxis coverage among bite victims (A) Comparison of unweighted and weighted annual incidence of animal bite and dog bite per 1000 population. Lines at the top of each bar represent 95% CIs. (B) Annual weighted dog-bite incidence by age, gender, and type of residence per 1000 population. Lines at the top of each bar represent 95% CIs. (C) Proportion of people bitten by dogs receiving ARV for at least one dose, three doses, and after completion of the prescribed schedule. (D) Proportion of people bitten by dogs who received RIG for various types of wounds. ARV=anti-rabies vaccine. RIG=rabies immunoglobulin.

Of the 1576 dog bites reported, 614 (39.0%) were by pet dogs and 1281 (81.3%) were unprovoked attacks. 1144 (72.6%) dogs were available for monitoring of their health status for 10 days, of which 1066 (93.2%) were alive after 10 days, 31 (2.7%) died, and 22 (1.9%) were killed. Most bites resulted in a single bite wound, predominantly on the lower limbs, upper limbs, torso, and the head or face. The most common type of wound was abrasion, followed by laceration, puncture wound, and avulsion (table 2).

Of the 1576 bite victims, 849 (53.9%) reported washing their wounds with soap and water and 276 (17.5%) applying some locally available remedies, such as chilli and turmeric powder (table 3). 1317 (83.6%) sought medical care at health facilities, with most seeking care within 24 h (appendix p 6).

Among people who had been bitten by dogs, 1253 (79.5%) received at least one dose of ARV, 1043 (66.2%) received three doses, and 638 (40.5%) received all the prescribed doses (figure 2; table 3; appendix p 7). Of the 1253 individuals who received at least one dose of ARV, 615 (49.1%) did not complete their full course of vaccines (appendix p 7). Of those who took vaccination, 1121 (89.5%) obtained it from public health

facilities and 836 (66.7%) were administered intradermally (table 3). The most common reasons for not taking the vaccine among unvaccinated individuals were lack of awareness, no belief in vaccination, and a minority took treatment from a traditional healer (appendix p 7). On multivariable analysis, odds of non-receipt of ARV were higher among dog-bite victims residing in north, northeast, east, and west regions. Other factors associated with non-receipt of ARV were bite victims with abrasion, being bitten by pet dogs, and residence in kuccha houses (appendix pp 9–11).

Only 81 (5.1%) bite victims reported taking rabies immunoglobulin. Among the 546 individuals with laceration, puncture, or avulsion wounds, only 61 (11.2%) had received rabies immunoglobulin (figure 2; table 3), mostly from public health facilities at the district and subdistrict levels (table 3).

Nearly a fourth of households interviewed reported having a pet or domestic animal. Of these, 3714 (18.6%) households reported that they had a pet dog and around half of the pet dogs were vaccinated (appendix p 12).

69 (4.4%) of the 1576 dog bites were classified as bites by suspected rabid dogs (appendix p 13), with the annual incidence of around 2.0 per 10000 population.

Between 2017 and 2022, 1880 suspected rabid dogs were tested by six rabies laboratories, with 1407 (74.8%) testing positive for rabies on direct fluorescent antibody testing (appendix p 14).

Of the 69 bites by suspected rabid dogs, 42 (60.9%) were on legs, 19 (27.5%) were on arms, six (8.7%) were on torsos, and two (2.9%) were on the head or face. 54 (78.3%) bite victims received at least three doses of ARV.

On the basis of the aforementioned data, the probability of death when bitten by a suspected rabid dog estimated using the decision-tree model was around 2% (table 4).

Using the annual incidence of suspected rabid-dog bites and the probability of death when bitten by a suspected rabid dog, we estimated 5726 (95% UI 3967–7350) human rabies deaths occur in India annually (table 4). The sensitivity analysis estimated 3908 (2741–5016) annual deaths based on the average bite incidence over the past 3 years. Considering the upper and lower 95% CIs of the data on rabies positivity among suspected rabid dogs, the estimated deaths were 5879 (95% UI 4078–7560) for the upper limit and 5573 (3897–7165) for the lower limit of rabies positivity among suspected rabid dogs (appendix p 15).

Discussion

Our survey estimated the annual incidence of animal bites in India to be 6.6 per 1000 persons, translating to 9.1 million bites annually, mostly caused by dogs. Nearly, two-thirds of the dog-bite victims had taken at least three doses of ARV. We estimated around 5700 human rabies deaths occur annually in India.

A systematic review in 2021 reported crude annual dog-bite incidence to be between 0.26% and 2.5%, with most studies reporting an incidence higher than 2%.¹² The dog-bite incidence observed in our study was lower than earlier studies. Higher incidences in earlier studies could be due to their smaller geographical scope and absence of probability sampling design. Our survey was nationally representative, using probability sampling at all levels of selection, with adequate sample size from each region of India. Another possible reason for lower incidence could be due to COVID-19 movement restrictions. However, the 1-year recall period of animal-bite history in 14 of the 15 states was after COVID-19 movement restrictions were lifted (appendix p 16), suggesting that our estimates might not have been greatly influenced by the pandemic. In 2019, the National Rabies Control Programme (NRCP) reported 7.2 million animal bites based on facility reporting.¹³ Our estimates of 9 million bites is close to the NRCP data before COVID-19, considering 20% of bite victims do not attend health facilities and 10% seek care from private health facilities.

Earlier studies in India estimated around 20 000 human rabies deaths annually.^{14,11} Compared to these estimates, there has been a substantial decline, nearly 75%, in

	People bitten by dogs
Wound management (n=1576)	
Wash with water and soap	849 (53.9%)
Locally available application	276 (17.5%)
Applied antiseptic	284 (18.0%)
Locally available application (n=276)	
Chilli powder	109 (39.5%)
Turmeric powder	107 (38.8%)
Plant sap or coconut water	10 (3.6%)
Other (mud, coffee powder, cow dung, coin snuff, powdered tablet, oil, or salt)	50 (18.1%)
ARV coverage by doses (either intradermal or intramuscular)	
Zero doses	323 (20.5%)
One dose	84 (5.3%)
Two doses	126 (8.0%)
Three doses	318 (20.2%)
Four doses	591 (37.5%)
Five doses	134 (8.5%)
Type of health facility ARV was received (n=1253)	
Public health facility (primary health centre or community health centre district hospital or subdistrict hospital or Taluk hospital medical college hospital)	1121 (89.5%)
Private health facility	132 (10.5%)
Route of administration (n=1253)	
Intradermal	836 (66.7%)
Intramuscular	417 (33.3%)
Site of ARV vaccination (n=1253)	
Deltoid	1217 (97.1%)
Anterolateral thigh	11 (0.9%)
Gluteal	25 (2.0%)
Received rabies immunoglobulin (n=1576)	
Type of health facility rabies immunoglobulin was received (n=81)	
Public health facility (primary health centre or community health centre district hospital subdistrict hospital or Taluk hospital or medical college hospital)	73 (90.1%)
Private health facility	8 (9.9%)
Type of bite wound and rabies immunoglobulin receipt	
Avulsion (n=11)	3 (27.3%)
Puncture (n=179)	30 (16.8%)
Laceration (n=356)	28 (7.9%)
Abrasion (n=1030)	20 (1.9%)
ARV=anti-rabies vaccination.	

Table 3: Post-exposure prophylaxis coverage among dog-bite victims

deaths over the past two decades. Analysis of data from the National Health Profile also showed a significant decline in incidence of human rabies death from 2.36 to 0.41 per 10 million population between 2005 and 2020.¹⁴ The decrease in deaths could be due to increased coverage of PEP among dog-bite victims. A nationwide study conducted in 2003 indicated 30.5% coverage of ARV and 2% coverage of rabies immunoglobulin in dog-bite victims.⁵ In our study, ARV coverage was 66.2% and rabies immunoglobulin coverage was 11.2%. Availability of ARV free of charge at public health facilities and greater awareness regarding ARV could have contributed to the increased ARV coverage.^{7,15}

	1-year survey data	Data source
Number of suspected rabid dog bites	69	Dog bites identified in the community survey were classified using Tepsumethanon's six criteria
Probability of a suspected rabid dog being confirmed rabid on laboratory diagnosis (1407 of 1880)	Binomial distribution: $p=0.748$	Using data from six laboratories in 12 Indian states
Bite injury to the head or neck (point estimate)	0.0290	Community survey data
Bite injury to the hand or arm (point estimate)	0.2754	Community survey data
Bite injury to the torso (point estimate)	0.087	Community survey data
Bite injury to the leg or foot (point estimate)	0.6087	Community survey data
Probability of developing rabies following a bite injury to the head by a rabid dog	Triangular distribution: minimum 0.30, mode 0.45, and maximum 0.60	Cleaveland et al ⁹
Probability of developing rabies following a bite injury to the hand or arm by a rabid dog	Triangular distribution: minimum 0.15, mode 0.28, and maximum 0.40	Cleaveland et al, ⁹ Knobel et al, ¹¹ and Fèvre et al ⁹
Probability of developing rabies following a bite injury to the torso by a rabid dog	Triangular distribution: minimum 0.00, mode 0.05, and maximum 0.10	Cleaveland et al, ⁹ Knobel et al, ¹¹ and Fèvre et al ⁹
Probability of developing rabies following a bite injury to the leg or foot by a rabid dog	Triangular distribution: minimum 0.00, mode 0.05, and maximum 0.10	Cleaveland et al, ⁹ Knobel et al, ¹¹ and Fèvre et al ⁹
Probability of an individual receiving anti-rabies vaccination if bitten by a suspected rabid dog (point estimate)	0.7826	Community survey data
Probability of death (p_{death})	0.02031828	$P1 \times ((P2 \times P6) + (P3 \times P7) + (P4 \times P8) + (P5 \times P9)) \times (1 - P10)$
Incidence of suspected rabid-dog bite (annual incidence)	69/337 808	Number of suspected dog bites in the surveyed population
At-risk population (Q)	1379 750 000	Population of India (2022)
Number of human rabies deaths (95% uncertainty limit) in 1 year	5726 (3967–7350)	$(p_{death}) \times (\text{annual incidence}) \times (Q)$

Table 4: Deaths due to human rabies estimated using the decision-tree probability model

Given that rabies is a fatal disease, timely, complete, and appropriate PEP is crucial. 20.5% of bite victims did not receive any ARV, with absence of awareness being the most common reason for non-vaccination. Dog-bite victims with abrasion were more likely not to take ARV, possibly underestimating the severity of small wounds. Focused IEC messaging is needed to stress the importance of seeking medical care even for minor abrasion.

Two-thirds of the bite victims in our survey received at least three doses and only 40.5% completed their scheduled course of vaccination. Additionally, about half of the bite victims who received one dose did not complete their full course of vaccination. Currently, in India, the scheduled dates of vaccination are written on outpatient sheets and bite victims are advised to come on scheduled vaccination dates. However, there is no system to remind the bite victims about their due dates of vaccination. Implementing reminder mechanisms such as SMS or

automatic voice calls, and in-person visits or telephonic calls from health workers can reduce dropouts. NRCP could also consider adopting the WHO-recommended shortened 1-week abridged Cambodia regimen¹⁶ for better compliance. Our study indicated that 50–67% of bite-victims from northeastern and eastern states received ARV intramuscularly. NRCP should advocate intradermal administration of ARVs in the deltoid region.

Only 11.2% of eligible bite victims in our survey reported receipt of rabies immunoglobulin, mostly from tertiary-care facilities. Making rabies immunoglobulin widely available in public health facilities is challenging because of limited production and high cost.¹⁷ Alternatively, use of rabies monoclonal antibodies could be considered with adequate training of health staff for correct use. Around half of the bite victims washed their wounds with soap and water and 17.5% used irritants such as turmeric and chilli powder. Such practices might delay care seeking and must be discouraged through continuous awareness by IEC messaging to the community.

In India, data on suspected human rabies deaths and animal bites are collected through the Integrated Health Information Portal of Integrated Disease Surveillance Programme.¹⁸ Additionally, NRCP collects this information from public health facilities once per month.¹⁹ During 2012–22, NRCP reported 6644 suspected human rabies cases and deaths, whereas the National Health Profile reported 2863 cases between 2005 and 2020 on the basis of health management information system data.^{3,14} However, our study estimated around 5700 human rabies deaths every year. The under-reporting of rabies deaths could be caused by incomplete reporting from health institutions, deaths occurring outside hospital settings, patients not getting admitted to hospital, and misdiagnosis. One major barrier to reporting human rabies cases is that victims either rarely get or do not get admitted to hospital. Additionally, the majority of human rabies cases are suspected based on clinical symptoms and laboratory confirmation is not routinely feasible in all suspected cases. Human rabies is a notifiable disease in India,²⁰ and states need to leverage such legal framework to improve accurate and timely reporting of all suspected human rabies cases occurring within and outside health facilities.

Rabies surveillance in animals in India is limited to ad-hoc reporting of suspected rabid animals to selected veterinary laboratories,²¹ with no integration between human and animal surveillance systems. Integrated surveillance is crucial for guiding rabies control activities including monitoring health status of biting dogs if they involve several biting incidents or are suspected to be rabid, initiating field investigation to trace all human and animal contacts and assessing for vaccination. This one-health approach has been successful in Latin American and southeast Asian countries,^{2,22,23} and is vital for achieving the goal of Zero by 30 in India.

Mass vaccination of dogs is crucial for rabies elimination. Countries such as Mexico, Sri Lanka, and Bangladesh

have successfully reduced human rabies mortality through consistent annual mass dog vaccination campaigns.^{23–25} In India, dog vaccination is done alongside the Animal Birth Control (ABC) programme, but data on dog vaccination coverages are scarce, and implementation of the ABC programme is suboptimal.²⁶ High rabies positivity (75% during 2017–22) in suspected rabid dogs and low (50%) vaccination coverage in pet dogs underscore the need to improved dog vaccination coverage. Only a few smaller states, such as Goa and Sikkim, have achieved high dog-vaccination coverage resulting in reduced animal rabies positivity.^{27,28} Given India's diversity, national campaigns are challenging and resource intensive. In view of this, the national action plan recommends that states estimate dog population in their respective areas and identify hotspots for conducting mass dog vaccination for 3 years with a coverage of 70% in those identified areas by local bodies, such as municipalities and municipal corporations and Panchayats.

Our study had certain limitations. First, the bite history of the family members was provided by an adult household member available during the survey, potentially missing bites among absent members or milder bites. However, animal bite is a unique medical event and family members are more likely to remember it. Further, bite exposure for the last 1 year was taken for analysis and hence missing bite exposure might be minimal and not affect the overall estimate. Additionally, recalling the number of vaccine doses taken might have been challenging, potentially underestimating the coverage by doses. We compared the number of doses received among the bite victims who had a documented proof and those with recall. The coverage by receipt of at least three doses was similar between those having and not having documentation of vaccination (appendix p 17). Second, we used a decision tree and probability model to estimate deaths. This model does not account for delays in receipt of ARV and rabies immunoglobulin. Our survey identified five deaths due to human rabies in the past 5 years (incidence 0.29 per 100 000 per year), roughly translating into 5000 deaths nationally, aligning with the estimate of our model. Third, rabies-positivity data among suspected rabid dogs were obtained from six laboratories catering to 12 states, with the number of samples tested ranging between one to 1000 in different states. Some states, especially in the eastern region of India, were not testing dog samples for rabies or data were not made available. Given that most of the data reported were from reference laboratories, rabies positivity might be overestimated. To account for this, sensitivity analyses were done and the number of deaths estimated assuming different scenarios were similar. Further, an expert group meeting using the Delphi technique was held among the academic and field veterinarians of India and the animal positivity data were validated before using them in the model. Lastly, we used Tepsu-methanon's criteria to classify a bite as a suspected rabid-dog bite. Only dogs not monitored for 10 days or

those killed within 10 days were classified using this criterion. In such instances, bite victims had a narrow window period to observe the health status of the dog. This short time period could have led to the underestimation of the incidence of suspected rabid-dog bites. However, only 30% of bites were classified using Tepsu-methanon's criteria (appendix p 13) and the rest were decided on the basis of the living status of the dog.

In conclusion, although there was a significant decline in rabies deaths over the past two decades, the programme needs to intensify actions towards elimination of dog-mediated rabies by 2030. Adopting a focused one-health approach by integrating human and animal surveillance, ensuring timely administration of a full course of PEP, and accelerating dog vaccination across the country are crucial steps towards this goal.

Human Rabies Study Collaborators

Deepak B Patil, Raja Paramasivam, Rajesh R Pharande, Karam Pal Singh, Swapna Susan Abraham, Aparna Sujatha, Parthiban Manoharan, Dhinakar Raj Gopal, Sheela Saly T George, Kamran Zaman, Krishna Pandey, Neha Singh, Manda Punnam Chander, Sanghamitra Pati, Srashti Panwar, Vishnu B Menon, Bathini Guru Viswanath, Atefu Ali, Hemant Mahajan, Karulesh Jain, Rajni Kant, Reshmi Ann Varkey, Gandhamalla P Sabitha Rani, Chinnappan Prabakaran, Ramakrishnan Gopinath, Paparaju Ashok Kumar, Arunachalam Suresh, Duraisamy Augustine, Kanniyappan Ramu, Karunakaran Sathish Kumar, Afeeq Karumaeh, Aparna Kumarababu, Suresh Arumugam

Human Rabies Study Collaborators affiliations

College of Veterinary Sciences and Animal Husbandry, Kamdhenu University, Gujarat, India (D B Patil PhD); Madras Veterinary College, Chennai, Tamil Nadu, India (R Paramasivam PhD, P Manoharan PhD, D R Gopal PhD); Mumbai Veterinary College, Mumbai, India (R R Pharande PhD); ICAR-Indian Veterinary Research Institute, Bareilly, Uttar Pradesh, India (K P Singh PhD); State Institute for Animal Diseases, Thiruvananthapuram, Kerala, India (S S Abraham MVSc, A Sujatha MVSc, S S T George BVSc); Himalayan Institute Medical Sciences, Swami Rama Himalayan University, Dehradun, India (S Singh BHMS); ICMR-Regional Medical Research Centre, Gorakhpur, India (K Zaman MD, R Kant PhD); ICMR-Rajendra Memorial Research Institute of Medical Sciences, Patna, India (K Pandey MD, H Mahajan DrPH); Pt Jawahar Lal Nehru Memorial Medical College, Raipur, India (N Singh PhD, K Jain MD); Amrita Institute of Medical Sciences, Amrita Vishwa Vidyapeetham, Kochi, India (V B Menon MD, R A Varkey MPH); Agartala Government Medical College and MRHRU, Tripura, India (M P Chander PhD); ICMR-Regional Medical Research Centre, Bhubaneswar, India (S Pati MD, A Ali PGPHM); SHARE India, Hyderabad, India (B G Viswanath MD, G P S Rani MSW); ICMR-National Institute of Epidemiology, Chennai, India (C Prabakaran MA, R Gopinath MA, P Ashok Kumar BA, A Suresh MA, D Augustine MA, K Ramu MA, K S Kumar MA, A Karumaeh M.Sc, A Kumarababu MBA, S Arumugam PhD)

Contributors

JWVT, MVM, and SD did the literature search. JWVT, MVM, NSK, and SD did the study design. JWVT, MVM, NSK, SE, DSR, SAK, AKS, AM, BS, DG, MM, NV, PS, RKS, RS, SMK, SB, TR, VVY, AKS, AsS, BVT, DRP, MMR, MT, NVS, PG, PJ, SV, ShD, SKP, VP, AGAR, DC, AD, AS, CP, SJ, SKI, MR, did the data collection. JWVT, MVM, NSK, SD, SE, and RS did the data analysis. JWVT, MVM, NSK, SE, DSR, SAK, AKS, AM, BS, DG, MM, NV, PS, RKS, RS, SMK, SB, TR, VVY, AdS, AsD, BVT, KP, MMR, MT, NVS, PG, PJ, SV, ShD, SKP, VP, AGAR, DC, AD, AS, CP, DHA, MKS, SJ, and MR, did the project management. JWVT, MVM, NSK, SD, and SE did the data interpretation. JWVT, MVM, SD, and SE accessed and verified the data. JWVT, MVM, NSK, and SD wrote the first draft of the manuscript. All authors approved the final draft of the manuscript.

Declaration of interests

We declare no competing interests.

Data sharing

All key anonymised individual participant data collected during the study, along with a data dictionary, and excluding data on costs incurred for post-exposure prophylaxis (which will be part of a separate manuscript) are available upon request to the corresponding author, after approval of a proposal with a signed data access agreement.

Acknowledgments

We acknowledge the support provided by the Ministry of Health and Family Welfare, Government of India, the state and district health administrative officials, and primary health-care staff in the conduct of the survey. We especially thank the Division of Zoonosis, National Centre for Disease Control (NCDC), Directorate General of Health Services, Ministry of Health and Family Welfare, Government of India for the support provided to the study. We thank Mugunthan Narayanan Muthulakshmi, Mohamed Ansari Resavu Mydeen, Ala Saritha, and Nandhini Prabhakaran for the support in producing the software images and maps for the figures. We also acknowledge the following team members for their support in planning and the conduct of the survey: Madhu Monjuri Gohain, Arupjyoti Konwar, Simi Gogoi, Yajong Y Konyak, T Shalome Khongsai, Maya Kumari Gupta, Johan Doley, Manash Pratim Baruah, and Gunomoni Borah (Assam and Nagaland); Prabhat Kumar Niraj and Ujjwal Prakash (Bihar); faculties and postgraduate students of the Department of Community Medicine, Pandit Jawahar Lal Nehru Memorial Medical College, Raipur (Chhattisgarh); K P Santhosh, Rethesh Kumar R, Mufeeda Moidu, Shifa Habeeb, Melvi Johnson, Amina Rasheed, Anmol A, Taniya P K, Greeshma S Kumar, Rose Maria Siju, Hamna Aboobacker, Sruthi V, and Anjitha Ravi (Kerala); Vijaya Saini, Raghvendra Dubey, Mohit Sharma, Krishkant Garg, and Ramsharan Prajapati (Madhya Pradesh); Kailas Babari Gadekar, Manchindra Rohidas Karanjawane, Avamishkumar Surendrakant Pandey, Rahul Harishchandrd Jagtap, and Vishal Ratnakant Khond (Maharashtra); Aditya Pratap Bhanja, Sarbeswar Bhoi, Ekant Priya Sutar, and Kishore Jyoti Moharana (Odisha); Manpreet Kaur, Harpreet Kaur, Prabhdeep Kaur, Jitesh, Jaspreet Kaur, Madan Mohan, Yuvraj Narang, Saminderpal Kaur Dhillon, Prabhjot Kaur Kalsi, and Rakesh Pal (Punjab); T Ravichandran, D Murugan, and R Harikrishnan (Tamil Nadu); D Mamatha, Vandana Bathini, M D Faheem, and Chetanand Bolbunda (Telangana); Tripura Model Rural Health Research Unit, Sanghita Paul, Sanjoy Karmakar, Bratati Paul, Satrajita Roy, Kaushik Das, Birendra Mohan Murasing, Mina Debbarna, and Sujata Das (Tripura); Pankaj Saxena and Lilly Singh (Uttar Pradesh); and the Principal, Research, and Ethics Committee of the Himalayan Institute Medical Sciences, Swami Rama Himalayan University, Dehradun (Uttarakhand).

References

- Hampson K, Coudeville L, Lembo T, et al. Estimating the global burden of endemic canine rabies. *PLoS Negl Trop Dis* 2015; 9: e0003709.
- WHO, Food and Agriculture Organization of the United Nations, World Organization for Animal Health, Global Alliance for Rabies Control. Zero by 30: the global strategic plan to end human deaths from dog-mediated rabies by 2030. <https://www.who.int/publications/i/item/9789241513838> (accessed July 16, 2024).
- Ministry of Health and Family Welfare. Government of India. 2021. National action plan for dog mediated rabies elimination from India by 2030. <https://rabiesfreeindia.mohfw.gov.in/resources/uploads/PageContentPdf/169391359514.pdf> (accessed July 16, 2024).
- Sudarshan MK, Madhusudana SN, Mahendra BJ, et al. Assessing the burden of human rabies in India: results of a national multi-center epidemiological survey. *Int J Infect Dis* 2007; 11: 29–35.
- Sudarshan MK, Mahendra BJ, Madhusudana SN, et al. An epidemiological study of animal bites in India: results of a WHO sponsored national multi-centric rabies survey. *J Commun Dis* 2006; 38: 32–39.
- Sudarshan MK, Ashwath Narayana DH. Appraisal of surveillance of human rabies and animal bites in seven states of India. *Indian J Public Health* 2019; 63: S3–8.
- Sudarshan MK. Vision 2030: dog-mediated human rabies-free India. Action must begin now. *Indian J Public Health* 2017; 61: 1–2.
- WHO. WHO expert consultation on rabies: WHO TRS number 1012, third report. 2018. <https://www.who.int/publications/i/item/WHO-TRS-1012> (accessed July 16, 2024).
- Tepsumedhanon V, Wilde H, Meslin FX. Six criteria for rabies diagnosis in living dogs. *J Med Assoc Thai* 2005; 88: 419–22.
- Cleaveland S, Fèvre EM, Kaare M, Coleman PG. Estimating human rabies mortality in the United Republic of Tanzania from dog bite injuries. *Bull World Health Organ* 2002; 80: 304–10.
- Knobel DI, Cleaveland S, Coleman PG, et al. Re-evaluating the burden of rabies in Africa and Asia. *Bull World Health Organ* 2005; 83: 360–68.
- John D, Royal A, Bharti O. Burden of illness of dog-mediated rabies in India: a systematic review. *Clin Epidemiol Glob Health* 2021; 12: 100804.
- Government of India, Ministry of Fisheries, Animal Husbandry and Dairying, Department of Animal Husbandry and Dairying, Shrimati Pratima Mondal. Unstarred question no 416 to be answered on 5th December 2023. Attacks by stray dogs. <https://sansad.in/getFile/loksabhaquestions/annex/1714/AU416.pdf?source=pqals> (accessed July 16, 2024).
- Rubeshkumar P, Majella MG, Jahan N, Sakthivel M, Krishnamoorthy Y. Secular trends of rabies in India, 2005–2020: importance of surveillance and implications for elimination strategies. *Lancet Reg Health Southeast Asia* 2024; 20: 100322.
- Sharma S, Agarwal A, Khan A, Ingle G. Prevalence of dog bites in rural and urban slums of Delhi: a community-based study. *Ann Med Health Sci Res* 2016; 6: 115–19.
- WHO. Rabies vaccines: WHO position paper, April 2018. *Wkly Epidemiol Record* 2018; 16: 201–20.
- Gogtay N, Thatte U, Kshirsagar N, et al. Safety and pharmacokinetics of a human monoclonal antibody to rabies virus: a randomized, dose-escalation phase 1 study in adults. *Vaccine* 2012; 30: 7315–20.
- Toppo VG, Kishore J. National action plan for eliminating dog-mediated rabies from India: the way forward. *APCR J* 2023; 25: 38–42.
- National Rabies Control Program, National Center for Disease Control, Directorate General of Health Services, Ministry of Health and Family Welfare, Government of India. Reporting formats NRCP. <https://rabiesfreeindia.mohfw.gov.in/downloads> (accessed July 16, 2024).
- National Center for Disease Control, Directorate General of Health Services, Ministry of Health and Family Welfare, Government of India. Centre for one health. <https://ncdc.mohfw.gov.in/centre-for-one-health/> (accessed July 16, 2024).
- Vanak AI, Panchamia N, Banerji I, Nair M. A passive “OneHealth” surveillance system to track canine rabies in urban India. *medRxiv* 2022. <https://doi.org/10.1101/2022.02.23.22271029> (preprint).
- Vigilato MAN, Cosivi O, Knobl T, Clavijo A, Silva HMT. Rabies update for Latin America and the Caribbean. *Emerg Infect Dis* 2013; 19: 678–79.
- Lionel Harischandra PA, Gunesekera A, Janakan N, Gongal G, Abela-Ridder B. Sri Lanka takes action towards a target of zero rabies death by 2020. *WHO South East Asia J Public Health* 2016; 5: 113–16.
- Vigilato MAN, Clavijo A, Knobl T, et al. Progress towards eliminating canine rabies: policies and perspectives from Latin America and the Caribbean. *Philos Trans R Soc Lond B Biol Sci* 2013; 368: 20120143.
- WHO. The rabies elimination program of Bangladesh. A model for transformation and accomplishment. <https://www.who.int/news/item/10-04-2017-the-rabies-elimination-program-of-bangladesh> (accessed July 17, 2024).
- Animal Welfare Board of India. Government of India. Scheme for birth control and immunization of stray dogs and medical screening of and treatment of tuberculosis in urbanized monkeys of Delhi. <https://awbi.gov.in/uploads/regulations/163318462167abc%20guidelines.pdf> (accessed July 16, 2024).
- Byrnes H, Britton A, Bhutia T. Eliminating dog-mediated rabies in Sikkim, India: a 10-year pathway to success for the SARAH program. *Front Vet Sci* 2017; 4: 28.
- Gibson AD, Yale G, Corfmaj J, et al. Elimination of human rabies in Goa, India through an integrated One Health approach. *Nat Commun* 2022; 13: 2788.
- Fèvre EM, Kaboyo RW, Persson V, Edelsten M, Coleman PG, Cleaveland S. The epidemiology of animal bite injuries in Uganda and projections of the burden of rabies. *Trop Med Int Health* 2005; 10: 790–98.